



7-2017

A Latent Profile Analysis of Aggression and Victimization across Relationship Types Among Veterans Who Use Substances

RaeAnn E. Anderson

University of North Dakota, raeann.anderson@UND.edu

Erin E. Bonar

Maureen A. Walton

Jason E. Goldstick

Sheila A. M. Rauch

See next page for additional authors

Follow this and additional works at: <https://commons.und.edu/psych-fac>

 Part of the [Psychology Commons](#)

Recommended Citation

Anderson, RaeAnn E.; Bonar, Erin E.; Walton, Maureen A.; Goldstick, Jason E.; Rauch, Sheila A. M.; Epstein-Ngo, Quyen M.; and Chermack, Stephen T., "A Latent Profile Analysis of Aggression and Victimization across Relationship Types Among Veterans Who Use Substances" (2017). *Psychology Faculty Publications*. 8.

<https://commons.und.edu/psych-fac/8>

This Article is brought to you for free and open access by the Department of Psychology at UND Scholarly Commons. It has been accepted for inclusion in Psychology Faculty Publications by an authorized administrator of UND Scholarly Commons. For more information, please contact zeineb.yousif@library.und.edu.

Authors

RaeAnn E. Anderson, Erin E. Bonar, Maureen A. Walton, Jason E. Goldstick, Sheila A. M. Rauch, Quyen M. Epstein-Ngo, and Stephen T. Chermack

Tables: 5, Figures: 1

A Latent Profile Analysis of Aggression and Victimization across Relationship Types
Among Veterans Who Use Substances

RaeAnn E. Anderson, PhD^a, Maureen A. Walton, MPH, PhD^{b, e}, Erin E. Bonar, Ph.D.^b, Sheila A.
M. Rauch, PhD^{c, d}, Quyen M. Epstein-Ngo, Ph.D.^{e, f}, Stephen T. Chermack, PhD^{b, g}

also:

Corresponding Author:

RaeAnn E. Anderson, phone: 330-672-3158, fax: 330-672-3786

^a Department of Psychological Sciences, Kent State University, 144 Kent Hall, Kent, OH 44242

^b Department of Psychiatry, Addiction Center, University of Michigan, 4250 Plymouth Road, Ann Arbor, MI 48109

^c Department of Psychiatry and Behavioral Sciences, Emory University School of Medicine, 12 Executive Park Dr, NE, 2nd Floor, Atlanta, GA 30329

^d Mental Health Service Line, Atlanta VA Medical Center, (116C) 1670 Clairmont Rd, Decatur, GA 30033

^e Department of Emergency Medicine, Injury Center, University of Michigan, 2800 Plymouth Road, Suite B10-GG080, Ann Arbor, Michigan, 48109

^f Institute for Research on Women and Gender, University of Michigan, 1136 Lane Hall, Ann Arbor, Michigan, 48109

^g VA Ann Arbor Healthcare System, Psychiatry, (116C) 2215 Fuller Road, Ann Arbor, MI, US, 48105

Supported by Grants: Clinicaltrials.gov #NCT01337973 to Stephen T. Chermack

NIAAA K23AA022641 to Quyen Epstein-Ngo, NIDA K23DA036008 to Erin E. Bonar.

Abstract

Objective: This study examined patterns of violence victimization and aggression in both intimate partner and non-partner relationships among veterans, and used latent profile analysis to identify subtypes of violence involvement.

Methods: Participants were 841 substance use treatment-seeking veterans (94% male) from a large VA Medical Center who completed screening measures for a randomized controlled trial. Self-report measures were: substance use, legal problems, depression, and violence involvement.

Results: Past year violence involvement, including both intimate partner (IPV) and non-partner (NPV) were common in the sample; although NPV occurred at somewhat higher rates. When including either IPV or NPV aggression or victimization, over 48% reported involvement with physical violence, 31% with violence involving injury and 86% with psychological aggression. Latent profile analysis including both aggression and victimization in partner and non-partner relationships indicated a four profile solution: no-low violence (NLV, $n = 701$), predominantly IPV ($n = 35$), predominantly NPV ($n = 83$), and high general violence (HGV, $n = 22$).

Multinomial logistic regression analyses revealed that compared to the no-low violence group, the remaining three groups differed in demographics, depressive symptoms, alcohol and other drug use, and legal involvement. Individuals within each profile had different patterns of substance use and legal involvement with the participants with an HGV profile reporting the most legal problems.

Conclusions: IPV and NPV are relatively common among veterans seeking substance use treatment. Characteristics of violence and associated substance use, mental health, and legal difficulties may be useful in considering how to tailor substance use and mental health services.

A Latent Profile Analysis of Aggression and Victimization across Relationship Types Among Veterans Who Use Substances

Involvement with violence (i.e., aggression and/or victimization) is a significant public health concern affecting approximately 30% of Americans in their lifetimes and is associated with an array of psychosocial problems, including substance use (Resnick et al., 1997; Rhodes et al., 2009; Tjaden and Thoennes, 2000). Experiencing victimization is associated with an array of mental and physical health complaints and even decreased occupational functioning (Byrne, Resnick, Kilpatrick et al., 1999; Tjaden & Thoennes, 2000). Engaging in aggression is also associated with poorer health including substance abuse and risk of HIV (Dunkle, Jewkes, Nduna et al., 2006; Raj, Reed, Welles et al., 2008).

There is a strong relationship between substance use and violence (Murphy & Ting). For example, rates of intimate partner violence (IPV), including either aggression or victimization, often exceed 50% among patients in substance use disorder (SUD) treatment settings (Brown et al., 1998; Chermack et al., 2000; Schumm et al., 2009). The few studies that have also assessed violence in non-partner relationships (e.g., friends, strangers, acquaintances, etc.) among patients in SUD treatment also show similar rates of non-partner violence (NPV) (Chermack et al., 2000; Murray et al., 2008a). In fact, when examining past-year involvement in either IPV *or* NPV among SUD patients, greater than 70% reported physical aggression (Chermack et al., 2000). The purpose of the present study was to identify patterns of violence involvement in partner and non-partner relationships among SUD treatment seeking veterans.

Violence involvement in veterans

IPV and NPV are relatively common problems among veterans, with factors such as SUDs, posttraumatic stress disorder (PTSD), and more recent service era (the time period/conflict

during enlistment) related to higher rates of violence (Elbogen et al., 2014; Gierisch et al., 2013; LaMotte et al., 2014; Marshall et al., 2005). Rates of past year IPV involvement among veterans with SUDs have been estimated in the 42-54% range (Chermack et al., 2008; Teten et al., 2009). The elevation of violence involvement in veterans with SUDs is consistent with biospsychosocial theories postulating that historical factors (e.g., prior violence exposure), psychological characteristics (psychological symptoms), alcohol and/or drug use, and social/contextual factors impact the risk for violence involvement (Chermack et al., 2009; Beck et al., 2013).

Thus, despite progress in research regarding IPV treatment among veterans (for example, Marshall et al., 2005 and Taft et al., 2013), examination of aggression *and* victimization simultaneously, across both partner and non-partner relationships, is under-explored in this population. Veterans tend to have high rates of problems that are associated with aggression/victimization risk, including poorer physical health functioning, PTSD and exposure to violence, heavy alcohol use, etc. (Hoerster et al., 2012; Elbogen et al., 2014; Gierisch et al., 2013; LaMotte et al., 2014; Marshall et al., 2005). Given these previous findings, it is important to explore profiles of violence among veterans' with SUDs in order to assist with prevention and treating violence-related problems in this population.

Forms of violence

An additional limitation of prior work includes lack of measurement of multiple forms of violence (psychological, physical and violence involving injury). Specification of the form of violence is important because different forms of violence may have different correlates and clinical outcomes (Epstein-Ngo et al., 2014). Physical assault and injury-related violence are primarily differentiated by severity where physical assault comprises any behavior that is inflicted physically and intended to cause harm whereas injury is indicated by bone or tissue

damage, a need for medical attention, or continuing physical pain. Differentiating these forms of violence highlights meaningful clinical differences; when examining only physical assault, it is common to find that women are equally or more violent than men (Archer, 2000), but when focusing on injury, women are much more likely to be seriously injured than men (Archer, 2000). There is also greater understanding that psychological forms of abuse, causing harm through verbal (e.g., calling names) or nonverbal behavior (e.g., stomping out of the room), can cause psychological distress and are worthy of clinical attention (Coker et al., 2012). Thus, research that better characterizes the heterogeneity of violence involvement is needed to enhance interventions for veterans.

Subtypes: IPV vs. NPV

Prior research has focused on samples characterized by IPV and identified subtypes (e.g., family only aggressors, borderline/dysphoric, generally violent/antisocial) (Holtzworth-Munroe et al., 2003; Walsh et al., 2010), that by design, did not include individuals who engage in only NPV. In SUD settings, research has shown some differential correlates for IPV and NPV (Chermack et al., 2000, 2010; Epstein-Ngo et al., 2014), suggesting potential subtypes exist, but few studies have employed person centered approaches to include multiple forms of IPV and NPV. Notable exceptions examining violence subgroups inclusive of both IPV and NPV provide an interesting foundation for understanding potential violence profiles in SUD treatment populations. Chermack et al., (2009) examined violence related injury among those in SUD treatment, and compared those: injuring partners only, non-partners only, or both; those injuring non-partners or both had more severe mood and substance problems. Another study found that correlates of violence were similar across victimization and aggression, but were differentiated by relationship type (IPV, NPV, or both) and gender (Walton et al., 2007).

Current study

The current study fills a gap in the literature by providing descriptive information (rates, frequencies) of multiple forms of violence and including both aggression and victimization across relationship types, among veterans seeking SUD treatment. Specifically, we include a comprehensive measurement of multiple forms of both NPV and IPV aggression and victimization (psychological, physical, injury) and use Latent Profile Analysis (LPA) to empirically derive violence subtypes. We hypothesized that at least three different profiles would emerge (consistent with Holtzworth-Munroe et al., 2003 and others) but did not make specific hypotheses about the characteristics of these groups given the lack of research on NPV. We also expected that groups with greater violence involvement would report more severe clinical characteristics. We chose potential clinical correlates variables based on biosychosocial models of violence and prior research (Chermack et al., 2009; Taft et al., 2007; Teten et al., 2009).

Methods

Participants and Procedures

Participants were 841 veterans initiating treatment or receiving treatment after a break in care from the Ann Arbor Veterans Affairs (VA) Healthcare System from 2012-2015. Participants were recruited as part of screening for a randomized controlled trial (RCT) of an intervention for substance use and violence prevention (based on Chermack et al., 2015). The data presented here were obtained during screening for the RCT. Inclusion criteria for screening were: recent substance use and attendance in outpatient SUD or mental health clinics. Exclusion criteria for screening were: living outside the catchment area, inability to provide informed consent (e.g., cognitive difficulties), having a legal guardian, inability to speak/read English, current suicidal intent and plan, acute psychosis, and ongoing participation in another intervention study.

Participants were remunerated \$10.00 for completing screening measures which included several surveys and was designed to be completed in approximately 30 minutes. Participants were mostly Caucasian (71.5%), male (93.5%), non-partnered (69.6%), non-employed (75.9%), low income (average \$10,000 - \$19,999), middle-aged ($M = 48.2$, $SD = 13.3$). Nearly two thirds of the sample served in Vietnam (29.4%) or served in Afghanistan and/or Iraq (OEF/OIF) (29.8%). Nearly 18% ($n = 147$) of participants were mandated or recommended to treatment by the criminal justice system. Data were collected under the supervision and approval of the Institutional Review Board of Ann Arbor VA and the trial is registered on ClinicalTrials.gov (#NCT01337973).

Measures

Past-year violence. Past year IPV and NPV aggression and victimization was assessed using a shortened version of the Revised Conflict Tactics Scale (CTS2; Straus et al., 1996). The CTS2 uses paired items to assess both aggression (“I hit my partner”) and victimization (“My partner did this to me”) on a frequency scale ranging from “0 – Never” to “6 – 20+ times” over the past year. Following convention, we computed frequency variables for each scale using the midpoint anchor of each selected range; alphas were generally acceptable (Table 1). For the NPV assessment we modified the CTS2 to assess NPV, e.g., “I kicked someone other than my partner” (Chermack et al., 2009; Murray et al., 2008). The parent intervention study focused primarily on reducing physical aggression (including injury) as measured by the CTS2. Given the need to also measure psychological symptoms in our screening battery we included a shortened version of the psychological aggression subscale (4 items were eliminated). The assessment of interpersonal violence excluded violence experienced as a result of combat.

Current depression symptoms. The 9-item Patient Health Questionnaire (PHQ-9) was used to assess past two-week symptoms (from 0 = “not at all” to 3= “nearly every day”) of major depression consistent with the DSM-IV criteria. Items are summed to yield a depression severity score ranging from 0 to 27 (Kroenke and Spitzer, 2002); in this study, Cronbach’s alpha was .90.

Current PTSD symptoms. The PTSD Checklist (PCL) was used to assess PTSD symptoms. The PCL is a 17 item measure that ask participants to rate how frequently they have experienced each of the 17 PTSD symptoms corresponding to DSM-IV-TR criteria on a scale ranging from 1 (“not at all/never”) to 5 (“extremely/daily or almost daily”). Item 17, “feeling jumpy or easily startled” was accidentally not administered. For the multivariate model we used a PCL cut-off score of 50 to conservatively estimate probable PTSD.

Past 28-day substance use. Substance use was assessed using items from the University of Arkansas Substance Abuse Outcomes Module (SAOM; Smith et al., 2006). Participants indicated the number of days they drank any alcohol as well as the number of days they consumed five or more drinks (e.g., heavy episodic drinking, HED). Similarly, participants indicated the number of days that they used each of the following: marijuana, cocaine or crack, stimulants (prescribed to participant and non-prescribed), opiates (prescribed to participant and non-prescribed), sedatives (prescribed to participant and non-prescribed), and heroin.

Lifetime legal problems. Items from the legal subsection of the Addiction Severity Index (ASI) were administered (McLellan et al., 1992). Because we were interested in distinguishing between legal problems related to aggression or other behaviors, we used factor analysis and correlation to create two composite scales from ASI items. The non-aggressive offenses scale comprised 8 items (e.g., shoplifting, vandalism, drug offenses, burglary, etc.), and the aggressive offenses scale (assault, rape, homicide, etc.) comprised 5 items. Following our

analyses, the DUI item was retained individually as it did not covary with other items significantly. To standardize the distribution, z scores were computed.

Statistical Analysis

LPA is a person-centered statistical approach that groups participants based on a set of measured variables that are hypothesized to represent a set of unmeasured or latent variables. In this analysis, twelve indicator variables (continuously scored), were used to include the three forms of aggression and victimization (physical assault, injury, psychological) in both partner and non-partner relationships separately.

We conducted the LPA in an exploratory fashion using Mplus (Version 6.11). Following standard LPA approaches, a two profile model was examined first and profiles added to the model until no additional statistical differences are found (Collins & Lanza, 2010). We evaluated model fit using Bayesian Information Criteria (BIC) with lower values indicating better fit, and entropy with values closer to 1.0 indicating better fit. The means of indicator variables within the profiles were examined to characterize each profile. Finally, we evaluated the model with the Vuong-Lo-Mendell-Rubin test and the Lo-Mendell-Rubin Adjusted Likelihood Ratio test to ensure that the model selected best accounted for the data as compared to a more parsimonious model, as indicated by p values greater than .05 for each of these tests. Drawing from the available (although limited) literature, our study is likely well-powered indicating the validity of the small groups extracted and model stability (Nylund, Asparouhov & Muthén, 2007).

To examine clinical correlates by profile, participants were assigned to their most likely profile using the probabilities generated by the LPA. Average profile probabilities for mostly likely profile membership for each class were: 1.00 for profile 1 (no-low violence), .97 for profile 2 (predominantly IPV), .98 for profile 3 (predominantly NPV), and 1.00 for profile 4

(high general violence). Profile assignments were exported from MPlus to SPSS; multivariate analyses were conducted in SPSS. In terms of missing data, no variable exceeded 2% that was missing, and missing data listwise (31 missing cases) for the multinomial logistic regression was < 4%. Following, we did not conduct data imputation and utilized list-wise deletion.

Bivariate comparisons (i.e., chi-square tests, analysis of variance) were made to examine differences between the profiles on demographic characteristics and mental health correlates. Some demographic characteristics were highly intercorrelated (service era and age; living situation and marital status) and these variables were pared in final models. Finally, we conducted a multinomial logistic regression analysis to evaluate multivariable relationships between the mental health correlates identified and violence profiles while accounting for shared variance.

Results

Descriptive Information Regarding IPV and NPV

Table 1 provides details regarding measures of IPV and NPV aggression and victimization in the sample. Collapsing across all three forms of violence and including victimization and aggression, 75.6% of participants reported any NPV, 64.7% reported any IPV, with 86.3% reporting any violence across relationships. IPV victimization and aggression were highly intercorrelated, $r(841) = .89, p < .001$. This was also true for any form of NPV victimization or aggression, $r(841) = .76, p < .001$.

Considering physical assault only, rates were 33.8% (NPV) and 23.2% (IPV), with 42.3% reporting either IPV or NPV. IPV physical assault victimization and physical assault aggression were highly correlated, $r = .71$, as was physical assault for NPV victimization and aggression, $r = .80, p < .001$. Rates of injury-related aggression or victimization ranged from approximately

10% to 22%. IPV physical assault and injury aggression were highly intercorrelated, $r(841) = .85$, $p < .001$ as were IPV physical assault and injury victimization, $r = .74$. Results were extremely similar for NPV. Finally, when including either IPV or NPV, over 48% reported involvement with physical assault, 31% with injury-related violence, and 86% for psychological abuse.

Latent Profile Analysis Findings

The means for each indicator variable in each profile are statistically different between each profile at $p < .001$; that is, for each of the twelve violence indicator variables, each profile is statistically unique from each other. A four profile solution best fit the data (BIC = 50596.47; entropy = 0.99). The Vuong-Lo-Mendell-Rubin test ($p = .61$) and Lo-Mendell-Rubin Adjusted Likelihood ratio test ($p = .61$) indicated that the three profile solution was similar but had a higher BIC = 51713.07. The Vuong-Lo-Mendell-Rubin and Lo-Mendell-Rubin Adjusted Likelihood ratio tests indicated a five profile solution was not statistically different ($p = .43$) from the four profile model; thus the more parsimonious four profile solution was chosen. See Table 2 for fit indices.

Table 3 displays the profile means and standard deviations for each indicator variable included in the LPA; Figure 1 graphically displays these means. Notably, these figures represent estimated frequency counts for each type of behavior; these counts do not necessarily represent separate incidents as a single altercation may include multiple behaviors, for example, both name calling and punching, which would be counted separately.

No-low violence profile.

The largest and least violent profile ($n = 701$ or 83.4%), was characterized as the no-low violence profile (NLV), where participants reported low violence across all three violence forms

and both social domains. For participants in this profile, their relationships are characterized by infrequent physical (M 's = 1.5 IPV, 9.7 NPV) and psychological (M 's = 18.1 IPV, 7.4 NPV) violence and rare injury for both victimization and aggression. Among individuals with this profile the means for counts of victimization were very similar to those for counts of aggression in the IPV domain.

Predominantly IPV profile.

The second profile ($n = 35$, 4.2%) was characterized as the predominantly partner violence (PIPV) profile because these participants reported high levels of IPV, particularly victimization, compared to NPV (although the levels of NPV were still far greater than NPV levels in the no/low violence profile). Participants in this profile reported an average of 50.1 counts of IPV physical victimization in the past year and an average of 2.9 counts of NPV physical victimization. These participants reported asymmetric violence in the IPV variables, reporting much more victimization than aggression counts of psychological, physical, and injury related violence. Profile 2 participants also reported asymmetry in the NPV domain, but this pattern differed: greater physical aggression, equivalent injury-related violence, and greater physical victimization.

Predominantly NPV profile.

The next profile ($n = 83$, 9.9%) was characterized as the predominantly non-partner violence profile (PNPV) as these participants reported the opposite pattern with relatively high levels of NPV. Mean counts of NPV physical victimization was 13.3 versus 4.0 for IPV physical victimization. In this profile, participants reported generally symmetric levels of violence within each relationship type, with the exception of NPV physical violence where they reported engaging in more aggression than victimization.

High general violence profile.

The final profile ($n = 22$, 2.6%) was characterized as the high general violence (HGV) profile and reported moderate to high levels of IPV as well as high levels of NPV. Of note, the high general violence profile reported the highest level of injury-related violence, reporting means two to four times those of other profiles. Individuals with this violence profile also reported a greater number of NPV counts than IPV counts and the highest levels of violence in either relationship type compared to all other profiles. Participant with this profile reported high symmetry in the domain of IPV but less so in the NPV domain, with more aggression than victimization counts reported. Of note, there were no women in this profile.

Differences in Clinical Characteristics among the Profiles

Results of chi-square and ANOVA tests (shown in Table 4) revealed that individuals comprising the four profiles did not vary significantly by current employment status, income, or education level. However, there were significant profile differences on age, gender, service era, and marital status. In relation to mental health correlates, the groups differed by substance use (HED, marijuana, cocaine/crack, stimulants, sedatives/hypnotics), legal involvement, depression symptoms, and PTSD symptoms, but not heroin, or opioid use (Table 4).

Results of the multinomial logistic regression model are shown in Table 5. Only variables identified as significant in the bivariate analyses were retained for the regression analyses; due to high intercorrelation ($r = .76$) between age and service era status, only age was included in this model. The no-low violence profile was used as the reference group. Gender was excluded from this analysis because the perfect prediction of this variable on to profile 4 destabilized the estimates of other variables.

Compared to the no-low violence profile, the predominantly IPV profile was younger, used cocaine/crack more frequently, used prescription stimulants more frequently, and had more non-aggression related legal charges. In comparison to the no-low violence profile, the predominantly NPV profile was younger, engaged in HED more frequently, had more aggression-related legal charges, and higher PTSD symptoms. Finally, compared to the no-low violence profile, the high general violence profile was younger, engaged in HED more frequently, had more severe PTSD symptoms and more legal charges (both aggression and non-aggression).

Discussion

Violence involvement is associated with many deleterious consequences including poor mental and physical health (Resnick et al., 1997). This study is the first to provide comprehensive information on violence involvement across multiple forms and relationship types among a treatment-involved sample of veterans. We found higher rates of NPV involvement than IPV involvement; this is consistent with other research that has examined both domains of violence in other SUD treatment settings (Chermack et al., 2009). The rates of both IPV and NPV in this sample were somewhat lower than those found in community or justice-referred samples (Beck et al. 2013; Crane et al. 2014), likely due to the higher mean age (48 years) and lower number of married couples in the present sample. The common presence of IPV and NPV highlights the importance of comprehensive screening for violence involvement in VA treatment clinics. When both NPV and IPV victimization and aggression measures were considered, 48% of the sample was identified as being involved with physical violence, 31% with injury-related violence, and > 86% with psychological violence.

The present study is unique in applying an advanced quantitative methodology, latent profile analysis, to “subtype” or characterize violence involvement by incorporating all violence variables simultaneously. Such an approach is critical and necessary to more fully understand potential typologies of violence and associated treatment needs for veterans. Indeed, our results are markedly different from much past research in IPV subtyping, likely reflecting the sample of veterans seeking SUD treatment and our comprehensive measurement of violence (Holtzworth-Munroe & Stuart, 1994; Langhinrichsen-Rohling et al., 2012). We identified four profiles of veteran SUD treatment seekers who were characterized primarily by the relationship type and severity of violence involvement, the latter a key indicator in prior research (Ansara and Hindin, 2010; Beck et al., 2013). The severity of violence in the profiles spanned the entire range from relatively low severity in the no-low violence group, moderate severity in the predominantly IPV and NPV profiles, to severe levels in the high general violence profile. Two of the profiles were identified by asymmetry in violence victimization and aggression (predominantly IPV and predominantly NPV), while the other two profiles (no-low violence and high general violence) had similar involvement in victimization and aggression across IPV and NPV domains. Notably, this violence excludes combat exposure.

Interestingly, the form of violence (e.g., the presence of injury-related violence) was not a unique characteristic in any profile, which is contrary to other latent class/profile research examining IPV using only partner victimization data and including more women (Ansara & Hindin, 2010; Beck et al., 2013). This may be partially due to the very high intercorrelations between physical assault and injury ($r > .7$), but it remains interesting that psychological aggression/victimization did not differentiate profiles. Contrary to some characterizations of IPV, this study did not find support for the “intimate terrorist” model of IPV (i.e., a group with high

levels of partner aggression with low levels of partner victimization) (Johnson, 2005). This may be partially due to our more comprehensive approach to measuring violence and the type of sample (e.g., veterans with substance misuse seeking treatment as opposed to court ordered domestic violence samples; our sample was older and less likely to be married than other samples). However, it is also possible that our predominately male participants (particularly in the high general violence profile) may have been reluctant to admit severe violence to or from partners. Our results are consistent with studies of non-veteran samples have suggested that the most common type of partner violence is “mutual aggression and victimization” (Straus, 2015). Further research using data from dyads is recommended (Leonard et al., 2015), although married dyads may not encompass all IPV given that such violence can occur in dating relationships and in extra-marital relationships as well.

We used the LPA results to identify how clinical characteristics consistent with biopsychosocial models of violence were differentially associated with the empirically derived profiles. Demographically the high general violence profile, was markedly different from the other groups. This group was the youngest and was entirely made up of men. Compared to the no-low violence group, all three profiles were more likely to experience PTSD symptoms, use substances, and be involved with the legal system. Notably, we were unable to determine whether legal problems predated substance use or vice versa; 17.7% of our sample was mandated/recommended to treatment but there was no significant difference between the identified profiles on this variable, $\chi^2(841) = .70, p = .87$.

Specific patterns of substance use and legal issues varied between groups with the predominantly IPV profile having a higher percent of women and being more likely to use a

range of substances, and the predominantly NPV and high general violence profiles more likely to engage in HED. The distinction of the predominantly IPV profile in having a higher percent of women and using a range of substances indicates there may be some unique treatment needs for women veterans and/or IPV involvement. Consistent with hypotheses and prior research, profiles with greater violence involvement also had more severe clinical characteristics, experiencing more PTSD symptoms and using more substances (Shorey et al., 2012, 2014; Taft et al., 2007). In terms of findings for substance use and violence across different relationship types, our findings are similar to prior research showing the association of alcohol and specific drugs may be different for partner and non-partner violence (Chermack et al., 2009; Chermack et al., 2010; Epstein-Ngo et al., 2014). We were unable to examine trauma history which may be a differentiating factor, particularly whether participants were exposed to early trauma or family violence. Although we investigated some of the clinical correlates most common in SUD clinics, we recommend future research investigate modifiable, individual difference variables in order to inform tailored intervention programs. For example, we recommend future research on constructs like trait anxiety, distress tolerance, and coping skills, which have been shown in prior research to be related to clinical outcome (Lejuez et al., 2008; Bornovalova et al., 2012; Levin et al., 2007).

Based on the different violence profiles identified in this paper, different types of clinical intervention may be warranted. Given the cross-sectional nature of this data, we present the following suggestions as potential hypotheses for future research. Our findings are clinically significant in identifying differences in the severity and type of violence involvement among veterans. Specific to each profile, the predominantly IPV profile may benefit from an intervention that targets relationship violence [e.g., Behavioral Couples Therapy (BCT)]

(O'Farrell & Schein, 2000), Strength at Home (Taft et al., 2013)] and poly-substance use; especially given evidence that violence remits as substance use remits (Chermack et al., 2015; Schumm et al., 2009). The predominantly NPV and high general violence group may benefit more from interventions more focused on alcohol and general violence prevention as well as case management given their legal problems. Prior studies suggest more positive SUD treatment outcomes with close coordination with the legal system (Friedmann et al., 2009; Crane et al., 2014). For example, the VA Veteran's Justice Outreach Program (e.g., coordinates substance use and mental health services and legal issues) shows positive clinical and legal outcomes (Slattery et al., 2013).

The present study has a number of limitations. First, participant reports of IPV, NPV or substance use were not able to be corroborated. Prior studies have tended to show acceptable concordance of participant and collateral reports of violence and substance use, with a general tendency for participants to under-report their own aggression (Epstein-Ngo et al., 2014; LaMotte et al., 2014; Murray et al., 2008b; Panuzio et al., 2006; Tharp et al., 2014); thus, it may be that the relationships identified in the present study are under-estimated. Future research should examine models in partnered vs. non-partnered participants to examine how relationship status may affect profile. Given the novelty of our findings and statistical approach, replication of these findings is required and longitudinal research needed to examine intervention outcomes as well as how profile membership may change over time and social context. We recommend future research improve typologies by including sexual violence, which may be relatively common in IPV involved couples and may be a differentiating factor (Bagwell-Gray et al., 2015). Finally, the present sample was veterans seeking treatment, with a relatively small percentage of women participants.

To summarize, the present study provides important and novel information regarding violence involvement (aggression and victimization) among veterans in VA treatment, including the identification of potential violence subtypes. Overall, violence involvement was common and those involved with more severe levels of violence had more severe clinical characteristics (substance use, PTSD symptoms, legal problems). Future research is needed to explore the impact of current and promising new approaches targeting violence prevention [e.g., Substance Abuse-Domestic Violence Treatment (Easton et al., 2007) and motivational interviewing/cognitive behavioral therapy (Chermack et al., 2015), etc.], including those that incorporate partners/families (BCT, Strength at Home), coordinate with the legal system when indicated, and/or require only the individual participant but target both IPV and NPV (which may have broader feasibility in clinical settings).

References

- Ansara, D. L., and Hindin, M. J. (2010). Exploring gender differences in the patterns of intimate partner violence in Canada: A latent class approach. *Journal of Epidemiology and Community Health*, *64*(10), 849–54. doi: 10.1136/jech.2009.095208
- Archer, J. (2000). Sex differences in aggression between heterosexual partners: a meta-analytic review. *Psychological bulletin*, *126*(5), 651.
- Bagwell-Gray, M. E., Messing, J. T., and Baldwin-White, A. (2015). Intimate partner sexual violence: A review of terms, definitions, and prevalence. *Trauma, Violence, & Abuse*, *16*(3), 316–335. doi: 10.1177/1524838014557290
- Beck, A., & Heinz, A. (2013). Alcohol-Related Aggression—Social and Neurobiological Factors. *Deutsches Ärzteblatt International*, *110*(42), 711–715.
<http://doi.org/10.3238/arztebl.2013.0711>
- Beck, C. J. A., Anderson, E. R., and O’Hara, K. L. (2013). Patterns of intimate partner violence in a large, epidemiological sample of divorcing couples. *Journal of Family Psychology*, *27*(5), 743–753. doi: 10.1037/a0034182
- Bornovalova, M. A., Gratz, K. L., Daughters, S. B., Hunt, E. D., & Lejuez, C. W. (2012). Initial RCT of a distress tolerance treatment for individuals with substance use disorders. *Drug and Alcohol Dependence*, *122*(1–2), 70–76. <http://doi.org/10.1016/j.drugalcdep.2011.09.012>
- Brown, T. G., Werk, A., Caplan, T., Shields, N., and Seraganian, P. (1998). The incidence and characteristics of violent men in substance abuse treatment. *Addictive Behaviors*, *23*(5), 573–586. doi: 10.1016/S0306-4603(98)00004-5
- Byrne, C. A., Resnick, H. S., Kilpatrick, D. G., Best, C. L., & Saunders, B. E. (1999). The

socioeconomic impact of interpersonal violence on women. *Journal of Consulting and Clinical Psychology*, 67(3), 362-6

Chermack, S. T., Bonar, E. E., Ilgen, M., Walton, M. A., Cunningham, R., Booth, B. M., and Blow, F. C. (2015). Developing an integrated violence prevention program for men and women in treatment for substance use disorders. *Journal of Interpersonal Violence*. doi: 10.1177/0886260515586369

Chermack, S. T., Murray, R. L., Walton, M. A., Booth, B. A., Wryobeck, J., and Blow, F. C. (2008). Partner aggression among men and women in substance use disorder treatment: correlates of psychological and physical aggression and injury. *Drug and Alcohol Dependence*, 98(1-2), 35–44. doi: 10.1016/j.drugalcdep.2008.04.010

Chermack, S. T., Fuller, B. E., and Blow, F. C. (2000). Predictors of expressed partner and non-partner violence among patients in substance abuse treatment. *Drug and Alcohol Dependence*, 58(1-2), 43–54. doi: doi.org/10.1016/S0376-8716(99)00067-8" da10.1016/S0376-8716(99)00067-8

Chermack, S. T., Grogan-Kaylor, A., Perron, B. E., Murray, R. L., De Chavez, P., and Walton, M. A. (2010). Violence among men and women in substance use disorder treatment: A multi-level event-based analysis. *Drug and Alcohol Dependence*, 112(3), 194–200. doi: 10.1016/j.drugalcdep.2010.06.005

Chermack, S. T., Murray, R. L., Walton, M. A., Booth, B. A., Wryobeck, J., & Blow, F. C. (2008). Partner aggression among men and women in substance use disorder treatment: correlates of psychological and physical aggression and injury. *Drug and Alcohol Dependence*, 98(1–2), 35–44. <http://doi.org/10.1016/j.drugalcdep.2008.04.010>

Chermack, S. T., Murray, R. L., Winters, J. J., Walton, M. A., Booth, B. M., and Blow, F. C.

(2009). Treatment needs of men and women with violence problems in substance use disorder treatment. *Substance Use & Misuse*, *44*(9-10), 1236–62. doi:

10.1080/10826080902960007

Coker, A. L., Smith, P. H., Bethea, L., King, M. R., & McKeown, R. E. (2000). Physical health consequences of physical and psychological intimate partner violence. *Archives of Family Medicine*, *9*(5), 451.

Collins, L. M., and Lanza, S. T. (2010). *Latent Class and Latent Transition Analysis: With Applications in the Social, Behavioral, and Health Sciences*. Wiley series in probability and statistics. Hoboken, N.J.: Wiley-Blackwell.

Crane, C. A., Schlauch, R. C., Hawes, S. W., Mandel, D. L., and Easton, C. J. (2014). Legal factors associated with change in alcohol use and partner violence among offenders.

Journal of Substance Abuse Treatment, *47*(2), 151–159. doi: 10.1016/j.jsat.2014.03.008

Dunkle, K. L., Jewkes, R. K., Nduna, M., Levin, J., Jama, N., Khuzwayo, N., Koss, M. P., et al.

(2006). Perpetration of partner violence and HIV risk behaviour among young men in the rural Eastern Cape, South Africa. *AIDS*, *20*(16), 2107-2114.

Easton, C. J., Mandel, D. L., Hunkele, K. A., Nich, C., Rounsaville, B. J., and Carroll, K. M.

(2007). A cognitive behavioral therapy for alcohol-dependent domestic violence offenders:

An integrated substance abuse-domestic violence treatment approach (SADV). *The*

American Journal on Addictions, *16*(1), 24–31. doi: 10.1080/10550490601077809

Elbogen, E. B., Johnson, S. C., Wagner, H. R., Sullivan, C., Taft, C. T., and Beckham, J. C.

(2014). Violent behaviour and post-traumatic stress disorder in US Iraq and Afghanistan veterans. *The British Journal of Psychiatry*, *204*(5), 368–375. doi:

10.1192/bjp.bp.113.134627

- Epstein-Ngo, Q. M., Walton, M. A., Sanborn, M., Kraus, S., Blow, F., Cunningham, R., and Chermack, S. T. (2014). Distal and proximal factors associated with aggression towards partners and non-partners among patients in substance abuse treatment. *Journal of Substance Abuse Treatment*, 47(4), 282–292. doi: 10.1016/j.jsat.2014.05.005
- Gierisch, J. M., Shapiro, A., Grant, N. N., King, H. A., McDuffie, J. R., and Williams, J. J. W. (2013). *Intimate partner violence: Prevalence among U.S. military veterans and active duty servicemembers and a review of intervention approaches*. Department of Veterans Affairs. Retrieved from <http://www.ncbi.nlm.nih.gov/books/PMH0066989/>
- Hoerster, K. D., Lehavot, K., Simpson, T., McFall, M., Reiber, G., & Nelson, K. M. (2012). Health and health behavior differences: US Military, veteran, and civilian men. *American Journal of Preventive Medicine*, 43(5), 483-489.
- Holtzworth-Munroe, A., Meehan, J. C., Herron, K., Rehman, U., and Stuart, G. L. (2003). Do subtypes of maritally violent men continue to differ over time? *Journal of Consulting and Clinical Psychology*, 71(4), 728–740. doi: 10.1037/0022-006X.71.4.728
- Holtzworth-Munroe, A., and Stuart, G. L. (1994). Typologies of male batterers: Three subtypes and the differences among them. *Psychological Bulletin*, 116(3), 476–497. doi: 10.1037/0033-2909.116.3.476
- Johnson, M. P. (2005). Domestic violence: it's not about gender—or is it? *Journal of Marriage and Family*, 67(5), 1126-1130.
- Kroenke, K., and Spitzer, R. L. (2002). The PHQ-9: A new depression diagnostic and severity measure. *Psychiatric Annals*, 32(9), 509–515.
- LaMotte, A. D., Taft, C. T., Weatherill, R. P., Scott, J. P., and Eckhardt, C. I. (2014). Examining

intimate partner aggression assessment among returning veterans and their partners.

Psychological Assessment, 26(1), 8–15. doi: 10.1037/a0034579

Langhinrichsen-Rohling, J., Misra, T. A., Selwyn, C., and Rohling, M. L. (2012). Rates of bidirectional versus unidirectional intimate partner violence across samples, sexual orientations, and race/ethnicities: A comprehensive review. *Partner Abuse*, 3(2), 199–230. doi: 10.1891/1946-6560.3.2.199

Lejuez, C. W., Zvolensky, M. J., Daughters, S. B., Bornovalova, M. A., Paulson, A., Tull, M. T., ... Otto, M. W. (2008). Anxiety sensitivity: A unique predictor of dropout among inner-city heroin and crack/cocaine users in residential substance use treatment. *Behaviour Research and Therapy*, 46(7), 811–818. <http://doi.org/10.1016/j.brat.2008.03.010>

Leonard, K. E., Winters, J. J., Kearns-Bodkin, J. N., Homish, G. G., and Kubiak, A. J. (2014). Dyadic patterns of intimate partner violence in early marriage. *Psychology of Violence*, 4(4), 384–398. doi: 10.1037/a0037483

Levin, C., Ilgen, M., & Moos, R. (2007). Avoidance coping strategies moderate the relationship between self-efficacy and 5-year alcohol treatment outcomes. *Psychology of Addictive behaviors*, 21(1), 108.

Marshall, A. D., Panuzio, J., and Taft, C. T. (2005). Intimate partner violence among military veterans and active duty servicemen. *Clinical Psychology Review*, 25(7), 862–76. doi: 10.1016/j.cpr.2005.05.009

Martin, A., Rief, W., Klaiberg, A., and Braehler, E. (2006). Validity of the Brief Patient Health Questionnaire Mood Scale (PHQ-9) in the general population. *General Hospital Psychiatry*, 28(1), 71–77. doi: 10.1016/j.genhosppsy.2005.07.003

- McLellan, A. T., Kushner, H., Metzger, D., Peters, R., Smith, I., Grissom, G., Pettinati, H., et al. (1992). The fifth edition of the Addiction Severity Index. *Journal of Substance Abuse Treatment, 9*(3), 199–213. doi: 10.1016/0740-5472(92)90062-S
- Murphy, C. M., & Ting, L. (2010). The effects of treatment for substance use problems on intimate partner violence: A review of empirical data. *Aggression and Violent Behavior, 15*(5), 325–333. <http://doi.org/10.1016/j.avb.2010.01.006>
- Murray, R. L., Chermack, S. T., Walton, M. A., Winters, J., Booth, B. M., and Blow, F. C. (2008a). Psychological aggression, physical aggression, and injury in nonpartner relationships among men and women in treatment for substance-use disorders. *Journal of Studies on Alcohol and Drugs, 69*(6), 896–905.
- Murray, R. L., Chermack, S. T., Walton, M. A., Winters, J., Booth, B. M., and Blow, F. C. (2008b). Psychological aggression, physical aggression, and injury in nonpartner relationships among men and women in treatment for substance-use disorders. *Journal of Studies on Alcohol and Drugs, 69*(6), 896–905. doi: 10.15288/jsad.2008.69.896
- Nylund, K. L., Asparouhov, T., & Muthén, B. O. (2007). Deciding on the number of classes in latent class analysis and growth mixture modeling: A Monte Carlo simulation study. *Structural equation modeling, 14*(4), 535-569.
- O'Farrell, T. J., and Schein, A. Z. (2000). Behavioral couples therapy for alcoholism and drug abuse. *Journal of Substance Abuse Treatment, 18*(1), 51–54.
- Panuzio, J., O'Farrell, T. J., Marshall, A. D., Murphy, C. M., Murphy, M., and Taft, C. T. (2006). Intimate partner aggression reporting concordance and correlates of agreement among men with alcohol use disorders and their remale partners. *Assessment, 13*(3), 266–

279. doi: 10.1177/1073191106287792

- Raj, A., Reed, E., Welles, S. L., Santana, M. C., & Silverman, J. G. (2008). Intimate partner violence perpetration, risky sexual behavior, and STI/HIV diagnosis among heterosexual African American men. *American Journal of Men's Health*, 2(3), 291-295.
- Resnick, H. S., Acierno, R., and Kilpatrick, D. G. (1997). Health impact of interpersonal violence: II. Medical and mental health outcomes. *Behavioral Medicine*, 23(2), 65–78. doi: 10.1080/08964289709596730
- Rhodes, K. V., Houry, D., Cerulli, C., Straus, H., Kaslow, N. J., and McNutt, L.-A. (2009). Intimate partner violence and comorbid mental health conditions among urban male patients. *Annals of Family Medicine*, 7(1), 47–55. doi: 10.1370/afm.936
- Schumm, J. A., O'Farrell, T. J., Murphy, C. M., and Fals-Stewart, W. (2009). Partner violence before and after couples-based alcoholism treatment for female alcoholic patients. *Journal of Consulting and Clinical Psychology*, 77(6), 1136–46. doi: 10.1037/a0017389
- Shorey, R. C., Elmquist, J., Ninnemann, A., Brasfield, H., Febres, J., Rothman, E. F., Schonbrun, Y. C., et al. (2012). The association between intimate partner violence perpetration, victimization, and mental health among women arrested for domestic violence. *Partner Abuse*, 3(1), 3–21. doi: 10.1891/1946-6560.3.1.3
- Shorey, R. C., Stuart, G. L., McNulty, J. K., and Moore, T. M. (2014). Acute alcohol use temporally increases the odds of male perpetrated dating violence: A 90-day diary analysis. *Addictive Behaviors*, 39(1), 365–368. doi: 10.1016/j.addbeh.2013.10.025
- Slattery, M., Dugger, M. T., Lamb, T. A., and Williams, L. (2013). Catch, treat, and release: Veteran treatment courts address the challenges of returning home. *Substance Use & Misuse*, 48(10), 922–932. doi: 10.3109/10826084.2013.797468

- Smith, G. R., Burnam, M. A., Mosley, C. L., Hollenberg, J. A., Mancino, M., and Grimes, W. (2006). Reliability and validity of the Substance Abuse Outcomes Module. *Psychiatric Services, 57*(10), 1452–1460. doi: 10.1176/appi.ps.57.10.1452
- Straus, M. A. (2015). Dyadic concordance and discordance in family violence: A powerful and practical approach to research and practice. *Aggression and Violent Behavior, 24*, 83–94. doi: 10.1016/j.avb.2015.04.011
- Taft, C. T., Kaloupek, D. G., Schumm, J. A., Marshall, A. D., Panuzio, J., King, D. W., and Keane, T. M. (2007). Posttraumatic stress disorder symptoms, physiological reactivity, alcohol problems, and aggression among military veterans. *Journal of Abnormal Psychology, 116*(3), 498–507.
- Taft, C. T., Macdonald, A., Monson, C. M., Walling, S. M., Resick, P. A., and Murphy, C. M. (2013). “Strength at Home” group intervention for military populations engaging in intimate partner violence: Pilot findings. *Journal of Family Violence, 28*(3), 225–231. doi: 10.1007/s10896-013-9496-y
- Teten, A. L., Schumacher, J. A., Bailey, S. D., and Kent, T. A. (2009). Male-to-female sexual aggression among Iraq, Afghanistan, and Vietnam veterans: Co-occurring substance abuse and intimate partner aggression. *Journal of Traumatic Stress, 22*(4), 307–311. doi: 10.1002/jts.20422
- Tharp, A. T., Sherman, M. D., Bowling, U., and Townsend, B. J. (2014). Intimate partner violence between male Iraq and Afghanistan veterans and their female partners who seek couples therapy. *Journal of Interpersonal Violence*. doi: 10.1177/0886260514564067
- Tjaden, P., and Thoennes, N. (2000). *Full report of the prevalence, incidence, and consequences of violence against women (NCJ 183781)*. Washington, DC.

Walsh, Z., Swogger, M. T., O'Connor, B. P., Chatav Schonbrun, Y., Shea, M. T., and Stuart, G.

L. (2010). Subtypes of partner violence perpetrators among male and female psychiatric patients. *Journal of Abnormal Psychology, 119*(3), 563–74. doi: 10.1037/a0019858

Walton, M. A., Cunningham, R. M., Chermack, S. T., Maio, R., Blow, F. C., & Weber, J. (2007).

Correlates of violence history among injured patients in an urban emergency department: gender, substance use, and depression. *Journal of Addictive Diseases, 26*(3), 61-75. doi: 10.1300/J069v26n03_07

Table 1

Descriptive Statistics for Violence Variables, n = 841

Violence Form & Social Domain	Prevalence: % of sample	Mean, SD	Non-Zero Median	Range
IPV-PV, $\alpha=.89$	26.6%	4.2(16.8)	4.0	0 – 225.0
IPV-IV $\alpha=.50$	12.5%	0.9(3.8)	4.0	0 – 40.0
IPV-PsyV $\alpha=.77$	61.0%	13.1(19.8)	11.0	0 – 100.0
IPV-PA $\alpha=.89$	23.2%	2.7(12.6)	3.0	0 – 250.0
IPV-IA $\alpha=.71$	9.5%	0.4(1.6)	2.0	0 – 58.0
IPV-PsyA $\alpha=.73$	62.2%	13.0(18.7)	11.0	0 – 100.0
NPV-PV $\alpha=.84$	30.1%	3.0(9.3)	4.0	0 – 110.0
NPV-IV $\alpha=.69$	21.9%	1.5(5.3)	3.0	0 – 60.0
NPV-PsyV $\alpha=.70$	68.3%	9.7(14.5)	7.0	0 – 83.0
NPV-PA $\alpha=.90$	33.8%	4.4(13.7)	4.0	0 – 162.0
NPV-IA $\alpha=.84$	18.5%	1.6(6.9)	3.0	0 – 108.0
NPV-PsyA $\alpha=.76$	74.9%	13.4(18.5)	9.0	0 – 100.0

Note. IPV = intimate partner violence, NPV = non-partner violence, PV = physical violence victimization, IV = injury victimization, PsyV = psychological victimization, PA = physical violence aggression, IA = injury aggression, PsyA = psychological aggression

Table 2
Fit indices for latent profile models

Number of Profiles in Model	BIC	SSA-BIC	VLMR LRT	LMR LRT	Entropy
3-profile	51713.07	51554.29	-26515.19	1635.36	.99
4-profile	50596.47	50396.40	-25688.17	1190.55	.99
5-profile	49570.77	49329.42	-25086.10	1100.68	.99

Note. AIC = Akaike information criterion, BIC = Bayesian information criterion, SSA-BIC = sample size adjusted BIC, VLMR = Vuong-Lo-Mendell-Rubin, LMR = Lo-Mendell-Rubin, LRT = likelihood ratio test

Table 3

Means and Standard Deviations by Violence Domain and Latent Profile

Form of Violence	Profile 1 – NLV N = 701, 83.4%	Profile 2 – PIPV N = 35, 4.2%	Profile 3 – PNPV N = 83, 9.9%	Profile 4 – HGV N = 22, 2.6%
	Intimate Partner			
Victimization				
Physical violence	0.92(2.97)	50.14(47.79)	4.04(11.88)	37.41(40.67)
Injury	0.14(0.74)	9.34(9.90)	1.41(4.73)	8.50(10.96)
Psychological	8.99(13.76)	59.86(24.61)	17.33(20.11)	52.91(28.32)
Aggression				
Physical violence	0.59(1.98)	19.97(24.50)	4.39(9.62)	34.36(56.47)
Injury	0.10(0.61)	3.71(6.68)	0.95(2.21)	6.95(13.68)
Psychological	9.09(14.05)	43.74(21.82)	23.42(23.40)	50.72(12.07)
	Non-partner			
Victimization				
Physical violence	0.56(1.66)	2.91(4.43)	13.27(11.43)	40.95(26.13)
Injury	0.28(1.33)	1.03(2.09)	6.16(5.25)	25.14(15.49)
Psychological	6.66(10.69)	16.00(16.05)	24.93(19.44)	40.41(20.36)
Aggression				
Physical violence	9.16(13.30)	18.89(21.35)	35.17(20.94)	58.27(26.05)
Injury	0.14(0.62)	1.14(2.38)	5.81(5.15)	33.27(24.44)
Psychological	0.71(1.72)	6.20(9.16)	17.25(11.76)	70.00(32.95)

Note. All variable are significantly different in each individual profile at $p < .001$

NLV = no-low violence, PNPV = predominantly non-partner violence, PIPV = predominantly intimate partner violence, HGV = high general violence.

Table 4

Profile Differences in Clinical Characteristics, N = 841

Variables	NLV (1) N = 701, 83.4%	PIPV (2) N = 35, 4.2%	PNPV (3) N = 83, 9.9%	HGV (4) N = 22, 2.6%	Statistical Test (omnibus)
Age (<i>M, SD</i>)	49.51(13.09) ^{abc}	42.37(12.90) _b	42.07(12.64) ^a	39.00(12.57) ^c	$F(3, 837) = 14.49, p < .001$
Gender (% women)	6.3%	20.0%	4.8%	0%	$\chi^2(3, 841) = 12.40, p = .006$
Marital Status (dichotomous, % married)	26.0%	17.1%	14.5%	9.1%	$\chi^2(3, 841) = 9.20, p = .03$
OEF/OIF/OND Era	26.7%	45.7%	45.8%	45.5%	$\chi^2(3, 841) = 20.20, p < .001$
Numbers of days HED (<i>M, SD</i>)	8.59 (11.26)	7.11(9.55)	11.60(11.77)	12.76(11.11)	$F(3, 828) = 2.85, p = .04$
Number of days marijuana (<i>M, SD</i>)	3.99(8.87) ^a	8.29(12.09) ^a	5.93(10.30)	5.59(10.14)	$F(3, 834) = 3.42, p = .02$
Number of days cocaine/crack (<i>M, SD</i>)	1.48(5.21) ^a	4.63(9.41) ^a	2.58(6.23)	3.82(8.83)	$F(3, 835) = 5.13, p = .002$
Number of days stimulants – prescribed (<i>M, SD</i>)	.51(3.53) ^a	2.63(8.51) ^a	.68(3.70)	2.05(7.01)	$F(3, 832) = 4.03, p = .007$
Number of days sedatives/hypnotics – not prescribed (<i>M, SD</i>)	.52(3.31) ^a	2.17(6.64) ^a	1.39(4.08)	1.36(6.40)	$F(3, 835) = 5.58, p = .01$
Legal – aggressive charges (ZM, ZSD)	-.18(2.46) ^{ab}	.07(2.13)	.67(3.02) ^a	3.01(6.93) ^b	$F(3, 830) = 11.74, p < .001$
Legal – non aggressive charges (ZM, ZSD)	-.14(2.07) ^a	.88(5.63)	.06(1.21)	2.84(6.65) ^a	$F(3, 829) = 11.90, p < .001$
Legal – DUI (ZM, ZSD)	-.03(1.37) ^a	-.20(.64)	.03(1.02)	1.26(3.29) ^a	$F(3, 830) = 6.30, p < .001$
Depression symptoms (<i>M, SD</i>)	10.48(6.92) ^{abc}	15.20(5.85) ^a	13.21(5.50) ^b	16.45(6.38) ^c	$F(3, 837) = 13.65, p < .001$
PTSD total score (<i>M, SD</i>)	16.38(.62) ^{abc}	53.91(14.75) ^a	54.89(12.87) _b	61.09(14.74) ^c	$F(3, 837) = 25.30, p < .001$

Note. OEF/OIF/OND = Operation Enduring Freedom/Operation Iraqi Freedom/Operation New Dawn, NLV = no-low violence, PNPV = predominantly non-partner violence, PIPV = predominantly intimate partner violence, HGV = high general violence. Z = Z score. HED = heavy episodic drinking, PTSD = posttraumatic stress disorder.

Superscripted pairs of letters indicate differences in group comparisons.

In the entire sample, the number of participants that reported any level of the following substances were: HED, $n = 458$; marijuana, $n = 263$; cocaine/crack, $n = 133$; stimulants – prescribed, $n = 36$; sedatives/hypnotics – not prescribed, $n = 59$.

Table 5

Multinomial Regression Analysis of Clinical Characteristic by Profile Compared to the No-Low Violence Profile

Variable	Profile 2 – PIPV Exp(B)[95% CI]	Profile 3 – PNPV Exp(B) [95% CI]	Profile 4 – HGV Exp(B) [95% CI]
Age	.96[.93, .99]	.95[.93, .97]	.91[.87, .96]
Marital status	.55[.21, 1.46]	.56[.28, 1.14]	.49[.10, 2.32]
# days HED	1.00[.96, 1.03]	1.03[1.01, 1.05]	1.05[1.00, 1.09]
# days marijuana	1.03[.99, 1.06]	1.01[.99, 1.04]	.98[.93, 1.04]
# days cocaine or crack	1.05[1.00, 1.09]	1.03[.99, 1.07]	1.02[.96, 1.10]
# days prescribed stimulants	1.06[1.00, 1.12]	1.00[.93, 1.07]	1.06[.99, 1.14]
# days NP sedatives or hypnotics	1.04[.97, 1.10]	1.05[.99, 1.10]	1.04[.95, 1.15]
Legal Charges – Aggression	1.00[.85, 1.17]	1.11[1.03, 1.20]	1.21[1.09, 1.34]
Legal Charges – Non-aggressive	1.10[1.00, 1.21]	1.01[.91, 1.13]	1.12[1.01, 1.24]
DUI	.80[.53, 1.21]	1.00[.82, 1.22]	1.07[.88, 1.30]
Depression Symptoms	1.06[.99, 1.14]	.98[.94, 1.03]	1.05[.96, 1.15]
PTSD	2.08[.84, 5.15]	4.42[2.31, 8.46]	5.12[1.28, 20.51]


Note. Bolded values indicate statistically significant differences at the $p < .05$ level or greater.

$R^2 = .18$ (Cox & Shell), $.26$ (Nagelkerke); Model $\chi^2(39) = 163.63$, $p < .001$

NP = non-prescribed. PNPV = predominantly non-partner violence, PIPV = predominantly intimate partner violence, HGV = high general violence. HED = heavy episodic drinking, PTSD = posttraumatic stress disorder.

Figure 1

Means by Violence Domain and Latent Profile



Note. IPV = intimate partner violence, NPV = non-partner violence, PV = physical violence victimization, IV = injury victimization, PsyV = psychological victimization, PA = physical violence aggression, IA = injury aggression, PsyA = psychological aggression, PNPV = predominantly NPV, PIPV = predominantly IPV, HGV = high general violence, NLV = no-low violence