



2018

Exercise in the Treatment of Major Depressive Disorder

Larissa Ann Haug
University of North Dakota

[How does access to this work benefit you? Let us know!](#)

Follow this and additional works at: <https://commons.und.edu/pas-grad-posters>



Part of the [Mental and Social Health Commons](#)

Recommended Citation

Haug, Larissa Ann, "Exercise in the Treatment of Major Depressive Disorder" (2018). *Physician Assistant Scholarly Project Posters*. 11.

<https://commons.und.edu/pas-grad-posters/11>

This Poster is brought to you for free and open access by the Department of Physician Studies at UND Scholarly Commons. It has been accepted for inclusion in Physician Assistant Scholarly Project Posters by an authorized administrator of UND Scholarly Commons. For more information, please contact und.common@library.und.edu.

Exercise in the Treatment of Major Depressive Disorder

Larissa Haug, PA-S

Department of Physician Assistant Studies, University of North Dakota School of Medicine & Health Sciences

Grand Forks, ND 58202-9037



Abstract

Objective

The purpose of this systematic literature review is to determine how exercise can be incorporated into the treatment of major depressive disorder (MDD) and its efficacy.

Methods


In this review, four databases were searched including SportDiscus, Pubmed, Cochrane Database of Systematic Reviews, and PsycInfo from October 1 to January 5, 2018. A variety of key terms were used when searching. Works chosen for review were published after the year 2000, were peer reviewed, and included randomized control trials (RCTs), pilot studies, systematic reviews, and meta-analyses. Sources that were excluded included those published prior to the year 2000, had poor study design, and included comorbid psychiatric conditions varying from MDD.

Results

For this review, 17 resources were selected. Much of the research presented shows evidence for the use of exercise in the treatment of MDD. However, most of the research points to exercise as more of a beneficial augmentation strategy for MDD versus a first line therapy involving pharmacotherapy with selective serotonin reuptake inhibitors (SSRIs) and psychotherapy in disease treatment.

Conclusion


Current research on the topic of exercise as therapy for MDD does show promise; however, more research still needs to be done in order to place exercise as an equivalent treatment to pharmacotherapy or psychotherapy.



The Hypothalamic-Pituitary-Cortisol System in Depression. Belmaker, R.H. & Agam, G. (2008). Major Depressive Disorder. *New England Journal of Medicine*, 358(55-68). doi: 10.1056/NEJMr073096

Research Question & Analysis

- What role exercise may have in the treatment of MDD?
 - Type, duration, and frequency of exercise most efficacious for MDD treatment will be assessed
 - Exercise will be compared to pharmacotherapy, psychotherapy, and exercise as augmentation to pharmacotherapy and/or psychotherapy
 - It is expected that exercise therapy will have benefits over SSRIs and psychotherapy but will not necessarily be superior to the latter two
- A systematic literature review will look at efficacy of the three therapies in regard to access, adherence, remission, and improvement of symptoms



Cooney, G. M., Dwan, K., Greig, C. A., Lawlor, D. A., Rimer, J., Waug, F. R., ... Mead, G. E. (2013). Exercise for depression. *Cochrane Database of Systematic Reviews*. doi:10.1002/14651858.CD004366.pub6

Literature Review

- Decreased hippocampal volume and function in MRI with increasing depressive episodes
- **Theme one: Type of Exercise, Intensity, and Dose for Therapeutic Benefit**
 - Neither aerobic or resistance training are superior to each other
 - Optimal therapeutic effect of exercise was 3 or more times per week for at least 30 minutes
 - Cochrane review of 39 studies showed that mixed aerobic and anaerobic exercise was most beneficial in MDD treatment and it took 13 to 36 sessions in order to see a large effect
 - Effect of exercise drops off after greater than 3 hours per week
 - Discontinuation of exercise causes worsening of depressive symptoms – not curative
- **Theme two: Exercise Compared to SSRIs and Psychotherapy**
 - Cochrane Review of 39 studies showed no significant difference comparing exercise to pharmacological and psychological treatment
 - Physical exercise was compared to usual care and placebo
 - Exercise compared to placebo showed a statistically significant improvement of depressive symptoms in mild to moderate MDD
- **Theme three: Exercise as Adjunctive Therapy for Depression**
 - Partial responders to SSRIs were assigned either a high or low dose exercise regimen; results showed significant improvement of depressive symptoms and quality of life in both groups
 - All participants were given 50 mg of sertraline and allowed to increase medication dose as needed; half of the participants were started on an exercise regimen as well; participants who were in the exercise group needed lower doses of sertraline but both groups had similar reduction in depressive symptoms
 - A study by Murri et al., randomized participants to sertraline plus non-progressive exercise, sertraline plus progressive exercise, and sertraline alone; showing a much higher remission rate for those in either exercise group compared to sertraline alone

Discussion

- One pitfall to many of the studies is that they excluded patients with MDD who were already living active life style and exercising; also, participants had to be willing to participate in an exercise regimen
- The study where participants got to choose their own intensity of exercise showed benefit, which is useful to know
- One study did show that the most effective dose for treatment in MDD is 30 minutes 3 times per week, which is not a very high number and may make exercise a more attractive treatment option for patients
- More adverse effects were seen in patients treated with pharmacotherapy versus exercise therapy
- Antidepressants such as sertraline are given a warning that they take weeks to months to take effect; therefore, starting patients on an exercise regimen at the same time as medication administration may result in a quicker improvement of depression symptoms

Applicability to Clinical Practice

- Exercise may play the best role as augmentation to psychotherapy or SSRIs pharmacotherapy as it relieves symptoms more rapidly than either of the two alone
- Starting patients on a mixed aerobic and resistance training program, 3 or more times per week is shown to be the most beneficial
- Patient empowerment is a major component to treating a patient; using exercise as a treatment gives the patients responsibility and control over their medical condition
- Concerns for using exercise as a therapy includes compliance, access, and ease of therapy; no doubt that taking a single pill daily is less time consuming than exercise
- Exercise has many other medical benefits including weight reduction, musculoskeletal strengthening, and cardiac conditioning
- More research needs to be done on the topic of exercise in the treatment of MDD

References

• Belmaker, R.H. & Agam, G. (2008). Major Depressive Disorder. *The New England Journal of Medicine*, 358, 55-68. doi:10.1056/NEJMr073096

• Bjornebekk, A., Mathe, A. A., & Brene, S. (2005). The antidepressant effect of running is associated with increased hippocampal cell proliferation. *International Journal of Neuropsychopharmacology*, 8, 357-368. doi:10.1017/S1461145705005122

• Cobb, J. A., Simpson, J., Mahajan, G. J., Overholser, J. C., Jurjus, G. J., Dieter, L., ... G., Stockmeier, C. A. (2012). Hippocampal volume and total cell numbers in major depressive disorder. *Journal of Psychiatric Research*, 47, 299-306.

• Cooney, G. M., Dwan, K., Greig, C. A., Lawlor, D. A., Rimer, J., Waug, F. R., ... Mead, G. E. (2013). Exercise for depression. *Cochrane Database of Systematic Reviews*. doi:10.1002/14651858.CD004366.pub6

• Dimeo, F., Bauer, M., Varahram, I., Proest, G., & Halter, U. (2001). Benefits from aerobic exercise in patients with major depression: a pilot study. *British Journal of Sports Medicine*, 35(2), 114-118. doi:10.1136/bjsm.35.2.114

• Doose, M., Ziegenbein, M., Hooe, O., Reim, D., Stengert, W., Hoffer, N., ... Sieberer, M. (2015). Self-selected intensity exercise in the treatment of major depression: A pragmatic RCT. *International Journal of Psychiatry in Clinical Practice*, 19(4), 266-276. doi:10.1080/13600567.2015.1082599

• Dynamid Plus. (2017). Major depressive disorder. Ipswich, MA: EBSCO Information Services. Retrieved October 30, 2017, Available from www.dynamid.com.

• Gartlehner, G., Gaynes, B. N., Amick, H. R., Asher, G. N., Morgan, L. C., Coker-Schwimer, E., ... Lohr, K. N. (2016). Comparative benefits and harms of antidepressant, psychological, complementary, and exercise treatments for major depression: An evidence report for a clinical practice guideline from the American College of Physicians. *Annals of Internal Medicine*, 164(5), 331-341. doi:10.7326/M15-1813

• Greer, T., Trombello, J., Rethorst, C., Carmody, J., Iha, M., Liao, A., ... Trivedi, M. (2016). Improvements in psychosocial functioning and health-related quality of life following exercise augmentation in patients with treatment resistant but nonremitted major depressive disorder: Results from the treat study. *Depression and Anxiety*, 33(9), 870-871. doi:10.1002/da.22521

• Halgren, M., Helgadóttir, B., Herring, M., Zeebani, Z., Lindfors, N., Kaldö, V., ... Forsell, Y. (2016). Exercise and internet-based cognitive-behavioral therapy for depression: Multicentre randomised controlled trial with 12-month follow-up. *The British Journal of Psychiatry*, 209(5), 414-420. doi:10.1192/bjp.bp.115.177576

• Hoffmann, B. M., Babayak, M. A., Craighead, W. E., Sherwood, A., Doraiswamy, M., Coons, M. J., & Blumenthal, J. A. (2011). Exercise and pharmacotherapy in patients with major depression: On year follow-up of the SMILE study. *Psychosomatic Medicine*, 73(2), 127-133. doi:10.1097/PSY.0b013e3182043a5

• Jonellsson, T., Lindvall, M., & Archer, T. (2014). Physical exercise intervention in depressive disorders: Meta-analysis and systematic review. *Scandinavian Journal of Medicine & Science in Sports*, 24(2), 259-272. doi:10.1111/sms.12050

• Marti, M. B., Amore, M., Menchetti, M., Toni, G., Neviani, F., Cerri, M., ... Zanetti, S. (2015). Physical exercise for late-life major depression. *The British Journal of Psychiatry*, 207(3), 235-242. doi:10.1192/bjp.bp.114.150516

• Nystrom, M., Neely, G., Hassmen, P., & Carlbring, P. (2015). Treating major depression with physical activity: A systematic overview with recommendations. *Cognitive Behavior Therapy*, 55(4), 341-352. doi:10.1080/16506073.2015.1015440

• Perce, B. W., O'Donnell, J. K., & Gaynes, B. N. (2012). The depression treatment cascade in primary care: A public health perspective. *Current Psychiatry Reports*, 14(4), 328-335. doi:10.1007/s11920-012-0274-y

• Siquiera, C. C., Valiengo, L. L., Cavalho, A. F., Santos-Silva, P. R., Missio, G., de Sousa, R. T., ... Machado-Vieira, R. (2016). Antidepressant efficacy of adjunctive aerobic activity and associated biomarkers in major depression: A 4 week, randomized, single-blind, controlled clinical trial. *PLoS One*, 11(5). doi:10.1371/journal.pone.0154195

• Stockmeier, C. A., Mahajan, G. J., Konick, L. C., Overholser, J. C., Jurjus, G. J., Meltzer, H. Y., ... Rajkowski, G. (2004). Cellular changes in the postmortem hippocampus in major depression. *Biological Psychiatry*, 56(9), 640-650.

• Yuan, T., Pais, F., Arias-Carrion, O., Rocha, N. B. F., Souza de Sa Filho, A., & Machado, S. (2015). Neural mechanisms of exercise: anti-depression, neurogenesis, and serotonin signaling. *CNS & Neurological Disorders - Drug Targets*, 14, 1307-1311.

Acknowledgements

I would like to thank my advisor Russell Kauffman and instructor Daryl Sieg for their patience and guidance with this large project. Another expression of gratitude must go to Dawn Hackman, our librarian, who kindly helped me with the initiation of my research and who patiently and promptly answered all my questions. I would also like to thank Marilyn G. Klug for her patience and time explaining the unfamiliar subject of statistics and for consequently improving my research skills. Finally, I would like to thank Tyler, Mom, Dad, Alayna, and Nicholas for the endless love and support through my physician assistant schooling.

Introduction

- Prevalence of MDD in the U.S. is 7%, lifetime prevalence is 16.6%
- MDD is a purely clinical diagnosis based on patient history and symptoms
- Hypotheses regarding the pathogenesis of depression:
 - An imbalance in the hormones serotonin, tryptophan, noradrenaline, and dopamine
 - Areas of the brain that show functional and structural abnormalities include anterior cingulate, orbitofrontal cortex, amygdala and basal ganglia, hippocampus, prefrontal cortex, ventral striatum, hypothalamic pituitary adrenal axis, and the nucleus accumbens

Statement of the Problem

- 1st line treatment for MDD involves psychotherapy or pharmacotherapy with SSRIs
 - Psychotherapy is difficult to access in rural areas with a shortage of trained counselors and psychiatrists
 - 40% of patients treated with SSRIs do not respond to therapy
 - Many patients treated with SSRIs complain of undesirable side effects
- Exercise is a potential benign, accessible, and affordable treatment option in mild to moderate MDD