5-20-2016

International normative 20 m shuttle run values from 1 142 026 children and youth representing 50 countries

Grant Tomkinson

University of North Dakota, grant.tomkinson@und.edu

Justin J. Lang

Mark S. Tremblay

Michael Dale

Allana G. LeBlanc

See next page for additional authors

[How does access to this work benefit you? Let us know!](#)

Follow this and additional works at: <https://commons.und.edu/ehb-fac>

Part of the [Health and Physical Education Commons](#)

Recommended Citation

Grant Tomkinson, Justin J. Lang, Mark S. Tremblay, et al.. "International normative 20 m shuttle run values from 1 142 026 children and youth representing 50 countries" (2016). *Education, Health & Behavior Studies Faculty Publications*. 26.

<https://commons.und.edu/ehb-fac/26>

This Article is brought to you for free and open access by the Department of Education, Health & Behavior Studies at UND Scholarly Commons. It has been accepted for inclusion in Education, Health & Behavior Studies Faculty Publications by an authorized administrator of UND Scholarly Commons. For more information, please contact und.commonson@library.und.edu.

Authors

Grant Tomkinson, Justin J. Lang, Mark S. Tremblay, Michael Dale, Allana G. LeBlanc, Kevin Belanger, Francisco B. Ortega, and Luc Léger

International normative 20 m shuttle run values from 1 142 026 children and youth representing 50 countries

Grant R Tomkinson, Justin J Lang, Mark S Tremblay, Michael Dale, Allana G LeBlanc, Kevin Belanger, Francisco B Ortega, Luc Léger

Abstract

Objective To develop sex-specific and age-specific international norms for the 20 m shuttle run test (20mSRT) in children and youth (aged 9–17 years), and to estimate the prevalence meeting the FITNESSGRAM criterion-referenced standards for healthy cardiorespiratory endurance (CRE).

Methods A systematic review was undertaken to identify papers explicitly reporting descriptive 20mSRT (with 1 min stages) data on children and youth since 1981. Data were included on apparently healthy (free from known disease/injury) 9–17 years old. Following standardisation to a common metric and for protocol differences, pseudo data were generated using Monte Carlo simulation, with population-weighted sex-specific and age-specific normative centiles generated using the Lambda Mu and Sigma (LMS) method. Sex-related and age-related differences were expressed as per cent and standardised differences in means. The prevalence with healthy CRE was estimated using the sex-specific and age-specific FITNESSGRAM criterion-referenced standards for $\dot{V}O_{2peak}$.

Results Norms were displayed as tabulated centiles and as smoothed centile curves for the 20mSRT using 4 common metrics (speed at the last completed stage, completed stages/minutes, laps and relative $\dot{V}O_{2peak}$). The final data set included 1 142 026 children and youth from 50 countries, extracted from 177 studies. Boys consistently outperformed girls at each age group (mean difference $\pm$ 95% CI: 0.86 $\pm$ 0.28 km/h or 0.79 $\pm$ 0.20 standardised units), with the magnitude of age-related increase larger for boys than for girls. A higher proportion of boys (mean $\pm$ 95% CI: 67 $\pm$ 14%) had healthy CRE than girls (mean $\pm$ 95% CI: 54 $\pm$ 17%), with the prevalence of healthy CRE decreasing systematically with age.

Conclusions This study provides the most comprehensive and up-to-date set of international sex-specific and age-specific 20mSRT norms for children and youth, which have utility for health and fitness screening, profiling, monitoring and surveillance.

Background

Cardiorespiratory endurance (CRE) is the ability to deliver oxygen to the muscles and to utilise it to generate energy to support muscle activity during exercise.^{1, 2} In adults, low CRE is strongly associated with cardiovascular disease and all-cause mortality and morbidity (independent of adiposity),^{3, 4} stroke,⁵ diabetes,⁶ mental health,⁷ health-related quality of life⁸ and cardiometabolic disease risk.^{6, 9} In children and youth, CRE is a weak-to-strong predictor of cardiovascular disease risk, cancer and mental health.^{10, 11} There is also evidence of an interaction between adiposity and CRE, suggesting that high levels of CRE may attenuate the deleterious effects of being overweight or obese in children and youth, the so-called ‘fat but fit’ phenotype.¹² Thus, CRE provides insight into the synergistic capabilities of several bodily systems and organs that are involved in the performance of physical activity and exercise, providing a strong and summative measure of health in children and youth.¹⁰ CRE also tracks moderately well from childhood to adulthood,^{13, 14} indicating that the measurement and surveillance of CRE in children provides insight into current and future population health statuses. Low fitness in childhood and adolescence is substantially linked with increased cardiometabolic disease risk,^{15, 16} obesity,¹⁷ reduced quality of life,¹⁸ poorer skeletal¹⁹ and mental health¹¹ in adulthood. In addition to health implications, CRE is an important

determinant of sporting success for many popular youth sports (eg, hockey, basketball, football (soccer), distance running, swimming, rugby).¹

The 20 m shuttle run test (20mSRT) is arguably the most popular field-based assessment and estimate of CRE in children and youth worldwide.^{20–23} It is an excellent tool for population-based surveillance and monitoring because it demonstrates strong-to-very strong test–retest reliability, and moderate-to-strong validity.²⁰ The 20mSRT also has excellent field-based utility due to its low cost, flexibility with testing locations (indoors, outdoors, smaller spaces) and its ability to test multiple individuals simultaneously with minimal equipment and personnel.^{20, 24, 25} In order to extend the utility of the 20mSRT as a surveillance instrument, there is a need for international norms, which, to date, have only been published for a single 20mSRT metric or for selected geographical regions,²⁰ including Europe,²⁶ North America²⁷ and Oceania.²⁸ Olds *et al*²¹ cumulated 20mSRT data on children and youth from 37 countries, representing six continents, suggesting that there is an international appetite for assessing CRE in children and youth. Harmonising reference values by creating international normative centiles for the 20mSRT in children and youth would provide opportunity for international surveillance and a means to compare CRE across geographic areas and time.

This study provides a 10-year update of the comprehensive 20mSRT reviews by Tomkinson *et al*²² and Olds *et al*,²¹ which were the first studies to describe a method to harmonise 20mSRT data by standardising for differences in test protocols and performance metrics. The primary aim was to develop the most comprehensive set of sex-specific and age-specific international normative centiles for CRE. The secondary aim was to estimate the proportion of children and youth meeting the FITNESSGRAM criterion-referenced standards for healthy CRE. These data will facilitate the identification of children and youth with very low CRE in order to set appropriate goals and promote positive health-related fitness behaviours, and conversely those with very high CRE, which may be important for sporting or athletic success.

Methods

Data sources

A systematic review of the scientific literature was registered (PROSPERO 2013:CRD42013003622) and completed to locate studies that reported descriptive 20mSRT data on 9–17 years old (see online supplement 1). Studies were identified up until October 2015 using the following bibliographic databases: MEDLINE (1946–2015), PsycINFO (1806–2014), EMBASE (1947–2014), SPORTDiscus (1949–2014) and Cochrane Central Register of Controlled Trials (2005–2014). The search strategy included the following terms: shuttle run*, OR beep test, OR multi-stage, OR aerobic, OR cardio*, OR endurance; with child*, adolescen*, pubescen*, boy, girl, young and youth as search term modifiers. All studies were extracted as text files, imported into Reference Manager (Thompson Reuters, San Francisco, California, USA), and assigned a unique reference identification number. Duplicate studies were first removed using Reference Manager with the remaining duplicates removed manually. Two independent reviewers screened all titles and abstracts for eligibility, with full-text copies obtained for all studies meeting initial screening criteria according to at least one reviewer. These two independent reviewers then examined all full-text articles and discrepancies were resolved by discussion and consensus. A third reviewer examined an article when the two reviewers were unable to reach consensus, with consensus reached for all included articles. Email contact with the corresponding authors of studies occurred when necessary, in order to provide clarification, to avoid ‘double counting’ previously reported data, and/or to request additional descriptive or raw data. The reference lists of all included studies were manually

reviewed to identify new studies. Reviewers contacted content experts to obtain grey literature. In addition, the extensive personal libraries of the study authors were examined for relevant studies.

Inclusion/exclusion criteria

Studies were included if they explicitly reported descriptive 20mSRT data (using the 1 min stage protocol; see Tomkinson *et al*²² for protocol variants) at the sex by age by country level. Study participants must have been apparently healthy (free from known disease or injury) 9–17 years old who were tested from 1981 onwards—the inception year of Léger's 20mSRT with 1 min stages. Studies were excluded if they reported descriptive data on: (1) other versions of the shuttle run (eg, the 15 m test or estimates of 20mSRT performance based on the 15 m test); (2) duplicate data published in another included study; or (3) on only special interest groups of children who were atypical of their source population (eg, elite athletes, physically or mentally retarded children). Figure 1 shows a flow chart of the included studies.

Figure 1
PRISMA flow chart outlining the identification of the included studies.

Data treatment and statistical analysis

All descriptive data were extracted into Excel (Microsoft Office 2010, USA) using a standardised data extraction table. The following descriptive data were extracted by one author and checked by another for accuracy: authors, country of testing, year of testing, sex, age, 20 m shuttle run protocol, 20 m shuttle run metric, sample size, mean, SD, median, sampling method and the sampling base. All data were examined for anomalies by running range checks and examining sex-specific and age-specific scattergrams. While only data on children and youth aged 9–17 years inclusive were retained for further analysis, individual study by sex by age by country by year groups were excluded when the sample size fell below 40 as the means and SDs for smaller samples were too labile. The final data set included 1896 study by sex by age by country by year groups.

The general procedure used to generate the sex-specific and age-specific normative centiles from extracted data is shown in figure 2. In most studies (76% or 135/177 studies), age was reported as age at last birthday; however, age was also reported in 7% (13/177) of studies as a span of years (maximum range: 3 years) and in 17% (29/177) as mean and SD years. In studies reporting age as a span of years, a new sample size was assigned to each study by sex by age by country group by dividing the reported sample size by the number of age groups (eg, a sample size of 162 was assigned to boys aged 10, 11 and 12 years in the study by Vandongen *et al*²⁹ which reported testing 486 boys aged 10–12 years). In studies that reported age as mean±SD years, Monte Carlo simulation was used to produce pseudo age data (using a random normal generator) based on reported means and SDs to estimate the sample size in each study by sex by age by country group.

Figure 2

Flow chart showing the methodological procedure used in this study. Results from studies were first expressed in the common metric of running speed (km/h) at the final completed stage and then corrected to Léger's original 1 min protocol according to which protocol they used (Léger, Eurofit or Queen's University of Belfast). Following the estimation of missing means (from reported medians) and SDs (from calculated CVs), poststratified population-weighted means and SDs were estimated for each sex by age group, with pseudo data then generated using Monte Carlo simulation. Smoothed centiles were then generated using the LMS method, with international normative 20mSRT values expressed in several different metrics (speed at the last completed stage, the number of completed full stages/minutes and relative $\dot{V}O_{2\text{peak}}$ (ml/kg/min)). 20mSRT, 20 m shuttle run test; CV, coefficient of variation; LMS, Lambda Mu and Sigma; and $\dot{V}O_{2\text{max}}$ peak oxygen uptake.

Testing year was recorded as the midpoint year of testing (eg, 2009.5 was recorded as the measurement year for a study that reported testing children in 2009) in 55% (98/177) of studies, with 34% (60/177) reporting a span of testing years and 11% (19/177) not reporting it at all. The midpoint year was recorded for studies reporting a span of testing years (eg, 2010.0 was recorded for a study reporting testing over the period 2009–2010), with 2 years prior to the publication year assumed for studies when it was not reported, which was the median difference for those studies in which the testing year was known.

To combine data from different studies, all 20mSRT data were standardised to a common metric and for protocol differences. To do this, we used an updated version of the standardisation procedure described in detail by Tomkinson *et al.*²² Figure 3 shows this updated standardisation procedure and summarises the steps used to express 20mSRT performances in the common metric of speed (km/h) at the last completed stage. All 20mSRT data were then standardised for protocol differences to Léger's original 1 min protocol,³⁰ which starts at a speed of 8.5 km/h and increases in speed by 0.5 km/h each minute.²²

Figure 3
Performance metric conversion flow chart adapted from Tomkinson *et al.*²² $\dot{V}O_{2peak}$, peak oxygen uptake.

As part of the modelling procedure used to generate sex-specific and age-specific normative centiles, means and SDs were required at the study by sex by age by country by year level. If no mean was available (1% or 2/177 studies), then mean values were estimated from the reported median values. This was done by first locating all studies reporting both median and mean values at the study by sex by age by country level, and second, by determining the best-fitting and most parsimonious linear and curvilinear (second and third order polynomials) regression models between median (predictor variable) and mean (response variable) speed values. Median and mean speed values were available for 418 study by sex by age by country groups, with the relationship nearly perfectly described by the following linear regression model: $\text{mean} = 0.9408 \times \text{median} + 0.6566$ (where r (95% CI) = 0.988 (0.985 to 0.990) and $SE = 0.128$). Furthermore, 5% (9/177) of studies did not report SD values. Missing SD values were estimated by first locating all studies reporting both means and SDs at the study by sex by age by country by year level; second, by calculating the corresponding coefficient of variation (CV) values; and third, by calculating the sample-weighted mean CVs for boys and girls separately. Mean and SD speed values were available

for 1585 study by sex by age by country groups, with sample-weighted mean CVs ($\pm 95\%$ CI) of $10.8 \pm 0.1\%$ and $9.2 \pm 0.1\%$ for boys and girls, respectively.

Sample-weighted means and SDs (the latter calculated from sample-weighted mean CVs) were then calculated at the sex by age by country level. While these data represent the best available 20mSRT data, in order to best generate internationally representative sex-specific and age-specific normative centiles and to correct for systematic bias associated with oversampling and undersampling, means and SDs were corrected using the poststratification population-weighting procedure described by Levy and Lemeshow.³¹ This procedure ensures that our population estimates reflect the underlying international age-specific and sex-specific country demographics. Thus, population estimates standardised to the mean measurement year of 2000 were extracted from the United Nations World Population Prospects report.³² Monte Carlo simulation was then used to create pseudo data using a random normal generator based on population-weighted means and SDs at the sex by age level. Monte Carlo simulation assumes that the distributions are approximately normal, which is true of the available raw 20mSRT speed data. Pseudo data sets were repeatedly generated until the calculated mean differed from the reported mean by $<0.5\%$, and the calculated SD differed from the reported SD by $<2.5\%$. These pseudo data sets were then used to generate sex-specific and age-specific normative centiles in LMSchartmaker Pro (V.2.43, The Institute of Child Health, London), which analyses data using the Lambda Mu and Sigma (LMS).³³ The LMS method fits smooth centile curves to reference data by summarising the changing distribution of three sex-specific and age-specific curves representing the skewness (L; expressed as a Box-Cox power), the median (M) and the CV (S). Using penalised likelihood, the curves can be fitted as cubic splines using non-linear regression, and the extent of smoothing required can be expressed in terms of smoothing parameters or equivalent degrees of freedom.³⁴ The effective degrees of freedom for 20mSRT speed were 1 (L curve), 4 (M curve) and 3 (S curve) for boys and 1 (L curve), 3 (M curve) and 3 (S curve) for girls. Normative centiles were also expressed in other common 20mSRT metrics, including the number of completed stages/minutes, the number of completed laps and relative peak oxygen uptake ($\dot{V}O_{2\text{peak}}$, mL/kg/min) values using the Léger *et al*³⁵ prediction equation:

$$\dot{V}O_{2\text{peak}}(\text{mL/kg/min}) = 31.025 + 3.238 \text{ speed} - 3.248 \text{ age} + 0.1536 \text{ speed} \times \text{age}$$

where speed is the running speed of the last completed stage (km/h) and age is age at last birthday. In a sample of Québec children and youth, this equation had a SE of estimate of 5.9 mL/kg/min or 12.1%.³⁵

The prevalence of children and youth (10–17 years old) with ‘healthy’ CRE was estimated using the new sex-specific and age-specific FITNESSGRAM criterion-referenced standards for $\dot{V}O_{2\text{peak}}$.^{36, 37} Differences in mean 20mSRT performance (km/h) between (1) age-matched boys and girls (eg, boys aged 10 years vs girls aged 10 years), and (2) sex-matched children of different ages (eg, boys aged 10 vs 11 years), were expressed as absolute and standardised differences. Positive differences indicated that 20mSRT performances for boys or older children were higher than those for girls or younger children. Standardised differences of 0.2, 0.5 and 0.8 were used as thresholds for small, moderate and large, respectively.³⁸

Results

The final data set included 1 142 026 children and youth aged 9–17 years (1896 study by sex by age by country by year groups extracted from 177 studies) from 50 countries tested over the period 1981–2014 (figure 4, see online supplement 2^{w39–w190}). These 50 countries represented six major geographical regions (Africa, Asia, Europe, Latin America and the Caribbean, Northern America and Oceania),³² including 33 high-income, 9 upper middle-income, 5 lower middle-income and 3 low-income economies.¹⁹¹ Norms are

presented as tabulated centiles from 5% to 95% for four common 20mSRT metrics in tables 1–4, with the smoothed centile curves for 20mSRT speed presented in figure 5 and the sex-specific and age-specific LMS values shown in online supplement 3.

Figure 4
World map representing the 50 countries (shown in black) for which 20mSRT data on 9–17 years old were available. 20mSRT, 20 m shuttle run test.

Figure 5
Smoothed centile curves (P10, P50 and P90) for the 20mSRT (speed (km/h) at the last completed stage) performance of (A) boys and (B) girls. 20mSRT, 20 m shuttle run test.

On average, approximately two-thirds of boys (mean±95% CI: 67±14%) and half of the girls (mean±95% CI: 54±17%) had healthy CRE, with the prevalence of healthy CRE decreasing by about 8% (boys) and 10% (girls) with every increasing year from age 10 to 17 years (figure 6).

Figure 6
Prevalence of 10–17 years old from 50 countries meeting the FITNESSGRAM Healthy Fitness Zone (V.10) thresholds for $\dot{V}O_{2\text{peak}}$ (mL/kg/min). $\dot{V}O_{2\text{peak}}$, peak oxygen uptake.

Boys consistently outperformed girls at each age group (mean difference $\pm$ 95% CI: 0.86 ± 0.28 km/h or 0.79 ± 0.20 standardised units), with the sex-related differences increasing with age from a small difference at age 9 years (difference in means $\pm$ 95% CI: 0.32 ± 0.01 km/h or 0.40 ± 0.01 standardised units) to a large difference at age 17 years (difference in means $\pm$ 95% CI: 1.46 ± 0.02 km/h or 1.20 ± 0.02 standardised units; figure 7A). From age 9 to 17 years, boys' performance improved at the rate of 0.17 km/h (or 0.15 standardised units) per year, with the largest rate of increase occurring at age 12 years (0.27 km/h or 0.23 standardised units). Girls' performance steadily improved at the rate of 0.03 km/h (or 0.03 standardised units) per year (figure 7B). These age-related changes were cumulatively large for boys and small for girls.

Figure 7
Standardised differences in mean 20mSRT performance (speed (km/h) at the last completed stage) between (A) age-matched boys and girls and (B) sex-matched children of different ages (anchored to age 17 years=0). Positive differences indicate that 20mSRT performances were higher for boys than for girls (top panel) or for older children than for younger children (bottom panel). The limits of the grey zone represent the thresholds for a large standardised difference (0.8 or -0.8). 20mSRT, 20 m shuttle run test.

Table 1

Twenty-metre shuttle run (speed (km/h) at the last complete stage) centiles by age and sex in 1 142 026 children and youth aged 9–17 years from 50 countries since 1981

Age (years)	P ₅	P ₁₀	P ₂₀	P ₃₀	P ₄₀	P ₅₀	P ₆₀	P ₇₀	P ₈₀	P ₉₀	P ₉₅
Boys											
9	8.63	8.94	9.31	9.58	9.82	10.03	10.25	10.48	10.75	11.13	11.44
10	8.61	8.95	9.35	9.65	9.90	10.13	10.36	10.61	10.90	11.31	11.64
11	8.60	8.97	9.41	9.72	10.00	10.25	10.50	10.77	11.09	11.53	11.89
12	8.64	9.05	9.53	9.89	10.19	10.47	10.75	11.05	11.40	11.89	12.29
13	8.74	9.18	9.72	10.10	10.43	10.73	11.04	11.37	11.75	12.29	12.72
14	8.85	9.32	9.88	10.29	10.64	10.96	11.29	11.64	12.04	12.61	13.08
15	8.94	9.42	10.01	10.43	10.79	11.13	11.47	11.83	12.25	12.84	13.32
16	9.01	9.51	10.11	10.55	10.92	11.27	11.62	11.99	12.42	13.03	13.53
17	9.08	9.60	10.22	10.67	11.05	11.41	11.77	12.16	12.60	13.23	13.74
Girls											
9	8.56	8.82	9.13	9.35	9.54	9.72	9.90	10.08	10.31	10.61	10.87
10	8.48	8.76	9.10	9.35	9.56	9.75	9.95	10.15	10.40	10.74	11.01
11	8.42	8.72	9.09	9.35	9.57	9.78	10.00	10.22	10.48	10.85	11.15
12	8.37	8.69	9.08	9.36	9.60	9.83	10.05	10.29	10.57	10.95	11.27
13	8.36	8.69	9.09	9.38	9.63	9.86	10.09	10.34	10.63	11.03	11.36
14	8.36	8.70	9.11	9.40	9.65	9.89	10.12	10.37	10.67	11.07	11.41
15	8.36	8.70	9.12	9.42	9.67	9.91	10.15	10.40	10.70	11.11	11.44
16	8.36	8.71	9.13	9.43	9.69	9.93	10.17	10.43	10.73	11.14	11.49
17	8.36	8.72	9.14	9.45	9.71	9.96	10.20	10.46	10.77	11.19	11.54

- The ages shown represent age at last birthday (eg, 9=9.00–9.99).

Table 2

Twenty-metre shuttle run (number of completed stages/minutes) centiles by age and sex in 1 142 026 children and youth aged 9–17 years from 50 countries since 1981

Age (years)	P ₅	P ₁₀	P ₂₀	P ₃₀	P ₄₀	P ₅₀	P ₆₀	P ₇₀	P ₈₀	P ₉₀	P ₉₅
Boys											
9	1.25	1.87	2.63	3.17	3.63	4.07	4.50	4.96	5.50	6.26	6.88
10	1.23	1.90	2.71	3.29	3.79	4.26	4.72	5.22	5.80	6.61	7.28
11	1.21	1.94	2.82	3.45	3.99	4.50	5.00	5.55	6.18	7.06	7.78
12	1.29	2.09	3.07	3.77	4.37	4.93	5.50	6.10	6.80	7.77	8.58
13	1.48	2.36	3.43	4.20	4.85	5.47	6.08	6.74	7.51	8.57	9.45
14	1.70	2.64	3.77	4.58	5.28	5.93	6.58	7.28	8.09	9.22	10.15
15	1.87	2.84	4.01	4.86	5.58	6.26	6.93	7.66	8.50	9.67	10.64
16	2.02	3.01	4.22	5.10	5.84	6.54	7.23	7.98	8.85	10.06	11.05
17	2.17	3.20	4.44	5.34	6.11	6.83	7.54	8.31	9.21	10.45	11.48
Girls											
9	1.12	1.63	2.25	2.70	3.08	3.43	3.79	4.17	4.61	5.23	5.74
10	0.97	1.53	2.21	2.70	3.11	3.50	3.89	4.31	4.80	5.47	6.03
11	0.83	1.44	2.17	2.70	3.15	3.57	3.99	4.44	4.96	5.69	6.29
12	0.74	1.39	2.17	2.73	3.21	3.65	4.10	4.57	5.13	5.90	6.54
13	0.72	1.38	2.19	2.77	3.26	3.72	4.19	4.68	5.25	6.05	6.71
14	0.72	1.40	2.21	2.80	3.31	3.78	4.24	4.75	5.33	6.14	6.81
15	0.73	1.41	2.24	2.83	3.34	3.82	4.29	4.80	5.39	6.21	6.89
16	0.73	1.42	2.26	2.87	3.38	3.86	4.34	4.86	5.46	6.29	6.97
17	0.73	1.44	2.29	2.90	3.43	3.92	4.40	4.93	5.54	6.38	7.08

Table 3

Twenty-metre shuttle run (number of laps) centiles by age and sex in 1 142 026 children and youth aged 9–17 years from 50 countries since 1981

Age (years)	P ₅	P ₁₀	P ₂₀	P ₃₀	P ₄₀	P ₅₀	P ₆₀	P ₇₀	P ₈₀	P ₉₀	P ₉₅
Boys											
9	9	14	20	24	28	32	36	40	45	52	58
10	9	14	21	25	29	33	37	42	47	55	62
11	9	15	22	27	31	36	40	45	51	60	67
12	9	16	24	29	34	39	45	50	57	67	75
13	11	18	26	33	39	44	50	56	64	75	84
14	13	20	29	36	43	48	55	62	70	81	92
15	14	22	31	39	45	52	58	66	74	86	97
16	15	23	33	41	48	54	61	69	78	91	102
17	16	25	35	43	50	57	64	72	81	95	107
Girls											
9	8	12	17	21	24	26	29	33	36	42	47
10	7	11	17	21	24	27	30	34	38	44	49
11	6	11	16	21	24	28	31	35	40	46	52
12	5	10	16	21	25	28	32	36	41	48	54
13	5	10	17	21	25	29	33	37	42	50	56
14	5	10	17	21	25	29	33	38	43	50	57
15	5	10	17	22	26	30	34	38	44	51	58
16	5	10	17	22	26	30	34	39	44	52	59
17	5	11	17	22	26	30	35	39	45	53	60

Table 4

Relative peak oxygen uptake ($\dot{V}O_{2peak}$, mL/kg/min) centiles by age and sex in 1 142 026 children and youth aged 9–17 years from 50 countries since 1981

Age (years)	P ₅	P ₁₀	P ₂₀	P ₃₀	P ₄₀	P ₅₀	P ₆₀	P ₇₀	P ₈₀	P ₉₀	P ₉₅
Boys											
9	41.7	43.1	44.8	46.1	47.1	48.1	49.1	50.2	51.5	53.2	54.6
10	39.7	41.3	43.2	44.6	45.8	46.9	48.0	49.2	50.6	52.5	54.1
11	37.7	39.5	41.7	43.2	44.6	45.8	47.0	48.4	49.9	52.1	53.9
12	36.0	38.0	40.5	42.3	43.8	45.2	46.7	48.2	50.0	52.4	54.5
13	34.6	36.9	39.7	41.7	43.4	45.0	46.6	48.3	50.3	53.1	55.4
14	33.3	35.8	38.8	41.0	42.9	44.6	46.4	48.3	50.5	53.5	56.0
15	31.8	34.5	37.8	40.1	42.1	44.0	45.9	47.9	50.2	53.4	56.1
16	30.4	33.2	36.6	39.1	41.3	43.2	45.2	47.3	49.8	53.3	56.1
17	28.9	32.0	35.6	38.2	40.5	42.6	44.7	46.9	49.5	53.2	56.2
Girls											
9	41.3	42.5	44.0	45.0	45.9	46.7	47.5	48.4	49.4	50.8	52.0
10	39.0	40.4	42.0	43.2	44.2	45.1	46.0	47.0	48.2	49.8	51.1
11	36.8	38.3	40.1	41.4	42.5	43.5	44.5	45.7	47.0	48.7	50.2
12	34.6	36.2	38.2	39.6	40.8	42.0	43.1	44.3	45.7	47.7	49.3
13	32.6	34.3	36.4	37.9	39.2	40.4	41.6	42.9	44.4	46.5	48.2
14	30.6	32.4	34.6	36.2	37.6	38.8	40.1	41.4	43.0	45.2	47.0
15	28.7	30.5	32.8	34.5	35.9	37.2	38.5	39.9	41.6	43.9	45.7
16	26.7	28.7	31.1	32.8	34.2	35.6	37.0	38.4	40.2	42.5	44.5
17	24.7	26.8	29.3	31.1	32.6	34.1	35.5	37.0	38.8	41.3	43.3

Discussion

This study systematically analysed 20mSRT data on 1 142 026 children and youth aged 9–17 years to generate the most comprehensive and up-to-date set of international sex-specific and age-specific norms for CRE. These international norms have utility for health and fitness screening, athlete profiling, and monitoring and surveillance in health, clinical, educational or sporting settings. They complement a

growing body of literature reporting national, regional and international growth centiles across a range of cardiometabolic disease risk factors, including adiposity (eg, body mass index^{192, 193} and waist circumference^{194–198}), blood pressure,^{199, 200} cholesterol,²⁰⁰ triglycerides,²⁰⁰ glucose²⁰⁰ and health-related fitness.^{28, 50, 126, 201–203}

Using a quintile framework, children in the bottom 20% can be classified as having ‘very low’ CRE; between the 20th and 40th centiles as having ‘low’ CRE; between the 40th and 60th centiles as having ‘moderate’ CRE; between the 60th and 80th centiles as having ‘high’ CRE; and above the 80th centile as having ‘very high’ CRE. Single measures of 20mSRT performance taken in health, clinical, educational or sporting settings can then be qualitatively interpreted using these quintile-based thresholds, with longitudinal changes (eg, due to growth or exercise training interventions) monitored by tracking changes against centile bands. For example, Armstrong *et al*¹ estimated that in children an appropriate 12-week CRE training programme will induce, on average, an 8–9% increase in $\dot{V}O_{2peak}$ independent of sex, age and maturation, equivalent to an increase of ~0.5 standardised units or ~20 centile points, which should be enough for a child to shift upwards from one quintile band to the next or above the relevant criterion-referenced standard for low cardiometabolic risk.²⁰⁴

While these norms are not criterion-referenced in that they do not indicate whether children have healthy CRE or low cardiometabolic risk, this study does provide an estimate of the prevalence with healthy CRE according to the new FITNESSGRAM (V.10) standards, which have been shown to discriminate with moderate accuracy between youth with and without metabolic syndrome.³⁷ There are currently no agreed on international criterion-referenced standards for CRE, and while we estimated prevalence using the US-based FITNESSGRAM standards, other national and regional standards have been published elsewhere.^{26, 205–211} It is important to note that the differences between published standards are (in some cases) substantial, with the new FITNESSGRAM (V.10) standards generally higher for girls and lower for boys relative to other standards,^{26, 205–211} meaning that our prevalence estimates would be substantially different if calculated using other standards. In contrast, $\dot{V}O_{2peak}$ can be estimated using different test protocols and prediction equations, and special care must be taken when comparing $\dot{V}O_{2peak}$ values with standards that were estimated using different test protocols and prediction equations.²¹² For example, we predicted $\dot{V}O_{2peak}$ from 20mSRT performance using the Léger *et al*³⁵ equation, whereas the new FITNESSGRAM standards were developed using predicted $\dot{V}O_{2peak}$ from a laboratory-based treadmill test.³⁷ To our knowledge, at least 17 prediction equations (from 10 studies of apparently healthy children and youth^{37, 213–221}) are currently available to estimate $\dot{V}O_{2peak}$ from 20mSRT performance, equations that differ in validity and can result in substantially different estimates of $\dot{V}O_{2peak}$ and hence the prevalence of healthy CRE. Future studies need to build on multinational efforts (eg, the HELENA study in Europe) in order to develop the most valid international normative-referenced and criterion-referenced standards for CRE.²²²

Using a cross-sectional approach, this study quantified the sex-related and age-related differences in 20mSRT speed, showing that boys consistently outperformed girls and experienced larger age-related changes. Given that longitudinal data are required to determine the true developmental patterns of CRE (because they control for within-participant changes in timing and tempo), and that larger age-related increases have been observed in children followed longitudinally than in those examined cross-sectionally,²²³ it is possible that our data underestimate the true developmental patterns. While the developmental patterns of children's $\dot{V}O_{2peak}$ has been well studied in non-representative longitudinal samples,^{224–226} other aspects of CRE (eg, mechanical efficiency, fractional utilisation, $\dot{V}O_2$ kinetics, etc) are less well understood, making it difficult to describe the mechanistic causes underlying the developmental patterns in 20mSRT performance. Given that relative $\dot{V}O_2$ and $\dot{V}O_{2peak}$ vary linearly with

speed and peak speed, changes in 20mSRT speed should therefore reflect changes in underlying $\dot{V}O_2$ (ie, the oxygen cost of the activity). Interestingly, however, this study observed decreases in boys' relative $\dot{V}O_{2peak}$ throughout childhood and adolescence, rather than the expected plateau.²²⁷ This unexpected developmental pattern may be due in part to $\dot{V}O_{2peak}$ prediction error or the fact that the developmental patterns were calculated using pooled cross-sectional data. In contrast, the premise of a plateau in boys' relative $\dot{V}O_{2peak}$ throughout childhood and adolescence is largely based on progressive treadmill or cycle testing of volunteer recruits who were athletically inclined, non-obese and motivated to exercise.^{226, 228} This unexpected finding in the developmental pattern in boys' $\dot{V}O_{2peak}$ might reflect current maturational development, or our use of a large, internationally representative sample of children and youth.

Strengths and limitations

This study brings together 20mSRT data from 177 studies in what is to date the largest international CRE database in the world. We conducted a systematic review, using a strict set of inclusion and exclusion criteria, applied rigorous data treatment procedures to systematically control for bias (eg, differences in test protocols and performance metrics), a poststratification population weighting procedure, a novel pseudo data method and the LMS method, to generate international sex-specific and age-specific smoothed centiles (across four performance metrics) for CRE. While it is not the first comprehensive study of children's 20mSRT performance, it updates the comprehensive reviews of Olds *et al*²¹ and Tomkinson *et al*²² by: (1) extending the data coverage from 2003 to 2014 through a rigorous systematic review process, (2) producing sex-specific and age-specific international normative centiles and (3) estimating the international prevalence of children and youth with healthy CRE.

Unfortunately, there are several sampling-related and methodology-related limitations to this study. First, it pooled data from studies that used different sampling methods which raises the issue of representativeness, with some studies using probability sampling (eg, stratified proportional, stratified random or cluster sampling) and others non-probability sampling (eg, convenience sampling; see online supplement 2). Second, differences in sampling base also exist across national, state/provincial and local/city/school levels. Third, differences in testing conditions (eg, environmental conditions such as climate or altitude, practice and running surfaces) and measurement errors (eg, calibration and type of equipment, methodological drift and diurnal variation) are inherent to any large data synthesis, but the very large number of data points captured should minimise these issues. Fourth, the vigorous nature of the 20mSRT may result in difficulties in test administration in, or exclusion of individuals with physical disabilities and diseases. The absence of data from these populations is likely to have inflated our normative values within the lower centile range.

The 20mSRT data were also collected at different times over the period 1981–2014, and given convincing evidence of international declines in children's CRE in recent decades,^{1, 23, 229} our norms may be biased. With a mean measurement year of 2000, and assuming an international decline of ~5% per decade since 1975,^{1, 23} our normative data may overestimate values in 2014 by ~7.5% (equivalent to ~0.65 standardised units or ~24 centile points) and represent a better health-related picture than what would be true today. However, without time trend data for all included countries and evidence of recent improvements in children's CRE in some included countries (eg, Australia,²⁸ Japan²²⁹ and Spain²³⁰), time-based corrections of our normative data are not recommended. Despite relying predominantly on data from high-income economies, there is no good evidence that 20mSRT performance is meaningfully related to a country's affluence or distribution of wealth,^{21, 231} so our norms are unlikely to be biased. Future studies need to examine CRE in low-income and middle-income economies and at multiple stages of the CRE transition. It must also be remembered that the 20mSRT is affected by factors other than

underlying construct CRE.^{20, 232} Validity data show that a moderate-to-large (35–70%) amount of the variance in 20mSRT performance is explained by the variance in underlying $\dot{V}O_{2peak}$,^{20, 232, 233} indicating that other physiological (eg, mechanical efficiency),²³⁴ fractional utilisation,²³⁵ $\dot{V}O_{2peak}$ kinetics,²³⁶ lactate threshold,²³⁷ anaerobic capacity,²³⁸ physical (eg, fat mass)²³⁹ and psychosocial factors (eg, motivation, effort and self-efficacy²⁰) also contribute.

Recommendations

Over the past few decades, the 20mSRT has been widely used to assess the CRE of children and youth, and yet data pooling is nearly impossible due to the difficulty with standardising performances (eg, because of differences in protocols, performance metrics, the way in which age is expressed, etc). To facilitate data pooling in the future, and to assist with the eventual update of 20mSRT norms, we make the following recommendations:

1. The test protocol used should be thoroughly and accurately reported;
2. Care should be taken to minimise and report factors that affect 20mSRT performance (eg, testing conditions and measurement errors);
3. Best practice should include 20mSRT results that are reported as the running speed (km/h) at the last completed stage;
4. Descriptive statistics (sample sizes, means and SDs) should be reported at the sex by age (at last birthday) level; and
5. The year(s) of testing should be reported;

Furthermore, because the 20mSRT is a maximal effort test, in order to ensure that a child has performed with ‘good effort’ (ie, they have tried very hard), perceptual (eg, ratings of perceived exertion) and/or physiological (eg, heart rate) effort should be measured in addition to performance effort.²⁴⁰ Any adverse events (or lack thereof) associated with maximal effort tests such as the 20mSRT should also be reported.²⁴¹

Conclusion

CRE is considered to be an excellent marker of current and future health. The 20mSRT is arguably the most popular measure of CRE because it is suitable for mass testing, is simple, cheap, easy, reliable, reasonably valid and is part of widely used health-related fitness test batteries (eg, Assessing Levels of Physical Activity and fitness (ALPHA),²⁴² Canadian Assessment of Physical Literacy (CAPL),²⁴³ Eurofit,²⁴⁴ FITNESSGRAM²⁴⁵ and even the PREFIT battery (Assessing FITness in PREscholers)).^{246, 247} Using a systematic review and analytical approach, this study used the best available 20mSRT data to: (1) provide the most comprehensive and up-to-date set of international sex-specific and age-specific norms for children and youth; and (2) estimate the prevalence with healthy CRE according to the FITNESSGRAM standards. These data have utility for health and sport promotion given that they help to identify children and youth with: (1) very low CRE in order to set appropriate fitness goals, monitor longitudinal changes and promote positive health-related fitness behaviours; and (2) very high CRE in the hope of recruiting them into elite sporting or athletic development programmes.

References

- 1 Armstrong N, Tomkinson GR, Ekelund U. Aerobic fitness and its relationship to sport, exercise training and habitual physical activity during youth. *Br J Sports Med* 2011;45:849–58.
- 2 Institute of Medicine. Health-related fitness measures for youth: cardiorespiratory endurance. In: Institute of Medicine, ed. *Fitness measures and health outcomes in youth*. Washington DC: The National Academy Press, 2012:111–51.
- 3 Blair SN, Kohl HW III, Paffenbarger RS Jr, *et al*. Physical fitness and all-cause mortality. A prospective study of healthy men and women. *JAMA* 1989;262:2395–401.
- 4 Kodama S, Saito K, Tanaka S, *et al*. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *JAMA* 2009;301:2024–35.
- 5 Lee CD, Blair SN. Cardiorespiratory fitness and stroke mortality in men. *Med Sci Sports Exerc* 2002;34:592–5.
- 6 Katzmarzyk P, Church TS, Janssen I, *et al*. Metabolic syndrome, obesity, and mortality: impact of cardiorespiratory fitness. *Diabetes Care* 2005;28:391–7.
- 7 Dishman RK, Washburn RA, Heath GW. *Physical activity epidemiology*. Champaign, IL: Human Kinetics, 2004:358–9.
- 8 Gillespie LD, Robertson MC, Gillespie WJ, *et al*. Interventions for preventing falls in older people living in the community. *Cochrane Database Syst Rev* 2009;(2): CD007146.
- 9 Katzmarzyk PT, Church TS, Blair SN. Cardiorespiratory fitness attenuates the effects of the metabolic syndrome on all-cause and cardiovascular disease mortality in men. *Arch Intern Med* 2004;164:1092–7.
- 10 Ortega FB, Ruiz JR, Castillo MJ, *et al*. Physical fitness in children and adolescence: a powerful marker of health. *Int J Obes* 2008;32:1–11.
- 11 Ruiz JR, Castro-Piñero J, Artero EG, *et al*. Predictive validity of health-related fitness in youth: a systematic review. *Br J Sports Med* 2009;43:909–23.
- 12 Eisenmann JC, Katzmarzyk PT, Perusse L. Aerobic fitness, body mass index, and CVD risk factors among adolescents: the Québec family study. *Int J Obes* 2005;29:1077–83.
- 13 Malina RM. Physical activity and fitness: pathways from childhood to adulthood. *Am J Hum Biol* 2001;13:162–72.
- 14 Ortega FB, Ruiz JR, Labayen I, *et al*. Role of socio-cultural factors on changes in fitness and adiposity in youth: a 6-year follow-up study. *Nutr Metab Cardiovasc Dis* 2013;23:883–90.
- 15 Dwyer T, Magnussen CG, Schmidt MD, *et al*. Decline in physical fitness from childhood to adulthood associated with increased obesity and insulin resistance in adults. *Diabetes Care* 2009;32:683–7.
- 16 Ortega FB, Silventoinen K, Tynelius P, *et al*. Muscular strength in male adolescents and premature death: cohort study of one million participants. *BMJ* 2012;345:e7279.

- 17 Ortega FB, Labayen I, Ruiz JR, *et al.* Improvements in fitness reduce the risk of becoming overweight across puberty. *Med Sci Sports Exerc* 2011;43:1891–7.
- 18 Morales PF, Sánchez-López M, Moya-Martinez M, *et al.* Health-related quality of life, obesity, and fitness in schoolchildren: the Cuenca study. *Qual Life Res* 2013;22:1515–23.
- 19 Moliner-Urdiales D, Ruiz JR, Ortega FB, *et al.* Association of physical activity with muscular strength and fat-free mass in adolescents: the HELENA study. *Eur J Appl Physiol* 2010;109:1119–27.
- 20 Tomkinson GR, Olds TS. Field tests of fitness. In: Armstrong N, van Mechelen W, eds. *Paediatric exercise science and medicine*. New York, NY: Oxford University Press, 2008:109–28.
- 21 Olds TS, Tomkinson GR, Léger L, *et al.* Worldwide variation in the performance of children and adolescents: an analysis of 109 studies of the 20-m shuttle run test in 37 countries. *J Sports Sci* 2006;24:1025–38.
- 22 Tomkinson GR, Léger LA, Olds TS, *et al.* Secular trends in the performance of children and adolescents (1980–2000): an analysis of 55 studies of the 20m shuttle run test in 11 countries. *Sports Med* 2003;33:285–300.
- 23 Tomkinson G, Olds T. Secular changes in pediatric aerobic fitness test performance: the global picture. *Med Sport Sci* 2007;50:46–66.
- 24 Ruiz JR, Silva G, Oliveira N, *et al.* Criterion related validity of the 20m shuttle run test in adolescents aged 13–19 years. *J Sports Sci* 2009;27:899–06.
- 25 Melo X, Santa-Clara H, Almeida JP, *et al.* Comparing several equations that predict peak VO₂ using the 20-m multistage-shuttle run-test in 8-10-year-old children. *Eur J Appl Physiol* 2011;111:839–49.
- 26 Silva G, Aires L, Mota J, *et al.* Normative and criterion-related standards for shuttle run performance in youth. *Pediatr Exerc Sci* 2012;24:157–69.
- 27 Carrel AL, Bowser J, White D, *et al.* Standardized childhood fitness percentiles derived from school-based testing. *J Pediatr* 2012;161:120–4.
- 28 Catley MJ, Tomkinson GR. Normative health-related fitness values for children: analysis of 85347 test results on 9-17-year-old Australians since 1985. *Br J Sports Med* 2013;47:98–108.
- 29 Vandongen R, Jenner DA, Thompson C, *et al.* A controlled evaluation of a fitness and nutrition intervention program on cardiovascular health in 10- to 12-year-old children. *Prev Med* 1995;24:9–22.
- 30 Léger L, Lambert J, Goulet A, *et al.* Capacité aérobie des Québécois de 6 à 17 ans —Test navette de 20 mètres avec paliers de 1 minute. *Can J Appl Sport Sci* 1984;9:64–9.
- 31 Levy PS, Lemeshow S. Stratification random sampling: further issues. In: Levy PS, Lemeshow S, eds. *Sampling of populations: methods and application*. Hoboken, NJ: John Wiley & Sons, Inc, 2008:143–88.
- 32 United Nations, Department of Economic and Social Affairs, Population Division (2015). *World Population Prospects: The 2015 Revision, Key Findings and Advance Tables* Working Paper No. ESA/P/WP.241.

- 33 Cole TJ, Green PJ. Smoothing reference centile curves: the LMS method and penalized likelihood. *Stats Med* 1992;11:1305–19.
- 34 Pan H, Cole T. *User's guide to LMSchartmaker*. UK: Medical Research Council, 2010:1–42.
- 35 Léger LA, Mercier D, Gadoury C, *et al*. The multistage 20 metre shuttle run test for aerobic fitness. *J Sports Sci* 1988;6:93–101.
- 36 Plowman SA, Meredith MD. *Fitnessgram/activitygram reference guide*. 4th edn. Dallas, TX: The Cooper Institute, 2013.
- 37 Welk GJ, Laurson KR, Eisenmann JC, *et al*. Development of youth aerobic-capacity standards using receiver operating characteristic curves. *Am J Prev Med* 2011;41: S111–16.
- 38 Cohen J. *Statistical power analysis for the behavioral sciences*. 2nd edn. New Jersey: Lawrence Erlbaum, 1988.
- 39 Secchi JD, García GC, Espana-Romero V, *et al*. Physical fitness and future cardiovascular risk in Argentine children and adolescents: an introduction to the ALPHA test battery. *Arch Argent Pediatr* 2014;112:132–40.
- 40 Hardy LL, King L, Espinel P, *et al*. *NSW Schools Physical Activity and Nutrition Survey (SPANS) 2010: full report*. Sydney: NSW Ministry of Health, 2010.
- 41 McIntyre S. *Trends in aerobic fitness from 1994 to 2009 in 10- and 11-year-old Australian children [thesis]*. Notre Dame, IN: University of Notre Dame, 2009.
- 42 Booth M, Okely AD, Denney-Wilson E, *et al*. *NSW Schools Physical Activity and Nutrition Survey (SPANS) 2004: full report*. Sydney: NSW Department of Health, 2006.
- 43 Lloyd KC, Antonas KN. Nutritional habits and fitness levels of schoolchildren. In: *Proceedings of the Nutrition Society of Australia—twenty-fourth annual scientific meeting, Fremantle, WA, 3-6 December*. Adelaide, SA: Nutrition Society of Australia, 2000:138.
- 44 Cooley D, McNaughton L. Aerobic fitness of Tasmanian secondary school children using the 20-m shuttle run test. *Percept Mot Skills* 1999;88:188–98.
- 45 Booth M, Macaskill P, McLellan L, *et al*. *NSW schools fitness and physical activity survey 1997*. Sydney: NSW Department of Education and Training, 1997.
- 46 Okely AD, Gray T, Cotton WG. Effect of an extended stay outdoor education program on aerobic fitness. In: Gray T, Hayllar B, eds. *Catalysts for change: proceedings from the 10th National Outdoor Education Conference*. Sydney, Australia, 1997:206–10.
- 47 Australian Council for Health, Physical Education and Recreation. *Australian fitness education award: user's manual and curriculum ideas*. Adelaide, SA: Australian Council for Health, Physical Education and Recreation, 1996.
- 48 Australian Sports Commission. *Norms for sport related fitness tests in Australian students aged 12-17 years*. Belconnen, ACT: Australian Sports Commission, 1994.
- 49 Jenner DA, Vandongen R, Beilin L. Relationships between blood pressure and measures of dietary energy intake, physical fitness, and physical activity in Australian children aged 11-12 years. *J Epidemiol Community Health* 1992;46:108–13.

- 50 Ortega FB, Artero EG, Ruiz JR, *et al.* Physical fitness levels among European adolescents: the HELENA study. *Br J Sports Med* 2011;45:20–9.
- 51 Vandendriessche JB, Vandenborgh BFR, Vaeyens R, *et al.* Variation in sport participation, fitness and motor coordination with socioeconomic status among Flemish children. *Pediatr Exerc Sci* 2012;24:113–28.
- 52 Seghers J, Rutten C. Clustering of multiple lifestyle behaviours and its relationship with weight status and cardiorespiratory fitness in a sample of Flemish 11- to 12-year olds. *Public Health Nutr* 2010;13:1838–46.
- 53 Verstraete SJ, Cardon GM, De Clercq DL, *et al.* A comprehensive physical activity promotion programme at elementary school: the effects on physical activity, physical fitness and psychosocial correlates of physical activity. *Public Health Nutr* 2007;10:477–84.
- 54 Cardon G, De Bourdeaudhuij I, De Clercq D, *et al.* Physical fitness, physical activity, and self-reported back and neck pain in elementary schoolchildren. *Pediatr Exerc Sci* 2004;16:147–57.
- 55 Heyters C, Marique T. *Le baromètre de la condition physique*. Bruxelles: Ministère de la Communauté française, Direction générale du sport, 2004.
- 56 Telama R, Naul R, Nupponen H, *et al.* *Physical fitness, sporting lifestyles and Olympic ideals: cross-cultural studies on youth sport in Europe*. Schorndorf, Germany: Verlag Karl Hofmann, 2002.
- 57 Baquet G, Berthoin S, Padovano C, *et al.* Effects d'un cycle de course de durée de type intermittent (court-court) sur la condition physique des adolescents. *Rev Educ Physique* 2000;40:51–60.
- 58 Lefèvre J, Bouckaert J, Duquet W, *et al.* De barometer van de fysieke fitheid van de Vlaamse jeugd 1997. *De resultaten. Sport (Blosso Brussel)* 1998;4:16–22.
- 59 Beunen G, Borms J, Vrijens J, *et al.* *Fysieke fitheid en sportbeoefening van de Vlaamse jeugd*. Volumen 1: Fysieke fitheid van de jeugd van 6 tot 18 jaar. Brussels: Blosso, 1991.
- 60 Poortmans J, Vlaeminck M, Collin M, *et al.* Estimation indirecte de la puissance aérobie maximale d'une population Bruxelloise masculine et féminine âgée de 6 à 23 ans. Comparaison avec une technique directe de la mesure de la consommation maximale d'oxygène. *J Physiol (Paris)* 1986;81:195–201.
- 61 Cazorla G, Gouthon P, Arémou, *et al.* *État de développement des capacités physiques des jeunes béninois 8-18 ans et plus. Rapport pour le Ministère Béninois de la Jeunesse et Sports et pour le Ministère Français de la Coopération*. Paris, France: Ministère Béninois de la Jeunesse et Sports et pour le Ministère Français de la Coopération, 1987.
- 62 Ribeiro RR, Santos KD, Carvalho WCG, *et al.* Aerobic fitness and biological and sociodemographic indicators in female school children. *Rev Bras Cineantropom Desempenho Hum* 2013;15:448–57.
- 63 Hobold E. *Indicadores de aptidão física relacionada à saúde de crianças e adolescentes do município de Marechal Cândido Rondon* [dissertation]. Paraná, Brasil: Universidade Federal de Santa Catarina, 2003.
- 64 Pieta S. *Estudio de la aptitud física de una muestra de la población escolar del estado de Paraná mediante la batería Eurofit* [dissertation]. León, Brasil: Universidad de León, 2000.

- 65 Veldhuizen S, Cairney J, Hay J, *et al.* Relative age effects in fitness testing in a general school sample: how relative are they? *J Sports Sci* 2014;33:109–15.
- 66 Voss C, Sandercock G, Higgins JW, *et al.* A cross-cultural comparison of body composition, physical fitness, and physical activity between regional samples of Canadian and English children and adolescents. *Can J Public Health* 2014;104: e245–50.
- 67 Leone M, Kalinova E, Comtois AS. *Global motor skill assessment from the UQAC-UQAM test battery: Canadian normative values by age and gender*. Québec: Université du Québec à Chicoutimi, 2011.
- 68 Reed KE, Warburton DER, Whitney CL, *et al.* Secular changes in shuttle-run performance: a 23-year retrospective comparison of 9- to 11-year-old children. *Pediatr Exerc Sci* 2006;18:364–73.
- 69 Massicotte D. *Partial curl-ups, push-ups and multistage 20 meter shuttle run, national norms for 6 to 17 year-olds. Final report submitted to Canadian Association for Health, Physical Education and Recreation (CAHPER) and Fitness and Amateur Sport Canada*. Montréal: University of Quebec at Montréal, 1990.
- 70 Garber MD, Sajuria M, Lobelo F. Geographical variation in health-related physical fitness and body composition among Chilean 8th graders: a nationally representative cross-sectional study. *PLoS ONE* 2014;9:e108053.
- 71 Kain J, Uauy R, Albala, *et al.* School-based obesity prevention in Chilean primary school children: methodology and evaluation of a controlled study. *Int J Obes* 2004;28:483–93.
- 72 Wong TW, Yu TS, Wang XR, *et al.* Predicted maximal oxygen uptake in normal Hong Kong Chinese schoolchildren and those with respiratory diseases. *Pediatr Pulmonol* 2001;31:126–32.
- 73 Wang P-G, Gong G, Wang SQ, *et al.* Relationship of body fat and cardiorespiratory fitness with cardiovascular risk in Chinese children. *PLoS ONE* 2011;6:e27896.
- 74 Cubides RC, Alarcón LGA, Galvi ARG. Diferencias en la actividad física y la condición física entre los escolares de secundaria de dos programas curriculares oficiales de Bogotá, Colombia. *Nutrición Hospitalaria* 2015;32:2228–34.
- 75 Ortega FJA. Estudio transversal de las cualidades funcionales de los escolares bogotanos: Valores de potencia aeróbica, potencia muscular, velocidad de desplazamiento y velocidad de reacción, de los siete a los dieciocho años. *Med Deporte Trab* 2013;32:1151–70.
- 76 Mojica GT, Poveda JG, Pinilla MI, *et al.* Sobrepeso, inactividad física y baja condición física en un colegio de Bogotá, Colombia. *Arch Latinoam Nutr* 2008;58:265–73.
- 77 Cazorla G, Dudal J, Dieu JL. *État de développement des capacités physiques des jeunes Ivoiriens 7-18 ans et plus. Rapport pour le Ministère Ivoirien de la Jeunesse et Sports et pour le Ministère Français de la Coopération*. Paris, France: Ministère Ivoirien de la Jeunesse et Sports et pour le Ministère Français de la Coopération, 1985.
- 78 Shahin A. Ying and yang of body composition assessment. *Chin J Integr Med* 2011;17:675–9.
- 79 Tinazci C, Emiroğlu O. Assessment of physical fitness levels, gender and age differences of rural and urban elementary school children. *Turkiye Klinikleri J Med Sci* 2010;30:1–7.

- 80 Gajda V. The applications of the UNIFITTEST battery at basic schools (age groups 7-14): summary report. Ostrava: PdF OU, 1994.
- 81 Nielsen GA, Andersen LB. The association between high blood pressure, physical fitness, and body mass index in adolescents. *Prev Med* 2003;36:229–34.
- 82 Cazorla G, Dudal J, Lefrancq L. *État de développement des capacités physiques des jeunes Djiboutiens 11-18 ans et plus. Rapport pour le Ministère Djiboutien de la Jeunesse et Sports et pour le Ministère Français de la Coopération*. Paris, France: Ministère Djiboutien de la Jeunesse et Sports et pour le Ministère Français de la Coopération, 1986.
- 83 Jürimäe T, Volbekiene V, Jürimäe J. Changes in Eurofit test performance of Estonian and Lithuanian children and adolescents (1992–2002). In: Tomkinson GR, Olds TS, eds. *Pediatric fitness: secular trends and geographic variability*. Basel: Karger, 2007:129–42.
- 84 Jürimäe T, Saar M. Self-perceived and actual indicators of motor abilities in children and adolescents. *Percet Mot Skills* 2003;97:862–6.
- 85 Jürimäe T, Volbekiene V. Eurofit test results in Estonian and Lithuanian 11 to 17-year-old children: a comparative study. *Eur J Phys Educ* 1998;3:178–84.
- 86 Raudsepp L, Jürimäe T. Relationships of physical activity and somatic characteristics with physical fitness and motor skill in prepubertal girls. *Am J Hum Biol* 1997;9:513–21.
- 87 Raudsepp L, Jürimäe T. Relationships between somatic variables, physical activity, fitness and fundamental motor skills in prepubertal boys. *Biol Sport* 1996;13:279–89.
- 88 Kull M, Jürimäe T. Using the Eurofit test battery in Estonian 16-18 years old adolescents. *Acta Commentationes Univ Tartuensis* 1994;967:49–52.
- 89 Palomäki S, Heikinaro-Johansson P, Huotari P. Cardiorespiratory performance and physical activity in normal weight and overweight Finnish adolescents from 2003 to 2010. *J Sports Sci* 2015;33:588–96.
- 90 Viljanen T, Taimela S, Kujala U. Koululaisten fyysinen aktiivisuus, kestävyyskunto ja ponnistuskorkeus. *Liikunta Tiede* 2000;37:23–6.
- 91 Baquet G, Twisk JWR, Kemper HCG, *et al.* Longitudinal follow-up of fitness during childhood: interaction with physical activity. *Am J Hum Biol* 2006;18:51–8.
- 92 Baquet G, Berthoin S, Gerbeaux M, *et al.* High-intensity aerobic training during a 10 week one-hour physical education cycle: effect on physical fitness of adolescents aged 11 to 16. *Int J Sports Exerc Med* 2001;22:295–300.
- 93 Baquet G, Berthoin S, Gerbeaux M, *et al.* Assessment of the maximal aerobic speed with the incremental running field tests in children. *Biol Sport* 1999;16:23–30.
- 94 Cazorla G, Portes A, James F. *Opération Martinique-Eval. Centre d'Evaluation Sport Santé, Fort de France (Martinique). Rapport pour L'Inspection d'Académie de la Martinique*. Fort de France, Martinique: L'Education d'Académie de la Martique, 1997.
- 95 Cazorla G. *Batterie France-Éval: Mesures, épreuves et barèmes: Évaluation des qualités physiques des jeunes Français d'âge scolaire: 7-11 ans. Rapport pour le Secrétariat d'Etat Auprès du*

- Premier Ministre Chargé de la Jeunesse et de Sports*. Paris: Ministère de la Jeunesse et de Sports, 1987.
- 96 Brunet J, Van Praagh E. *Batterie expérimentale de tests moteurs Eurofit: Rapport d'Activité de la Région Auvergne—1984-1985*. Clermont-Ferrand, France: Université de Clermont-Ferrand, 1985.
- 97 Tambalis KD, Panagiotakos DB, Psarra G, *et al*. Physical fitness normative values for 6-18-year-old Greek boys and girls, using the empirical distribution and the lambda, mu, and sigma statistical method. *Eur J Sport Sci* 2015;1–11.
- 98 Tambalis K, Panagiotakos D, Sidossis L. Greek children living in rural areas are heavier but fitter compared to their urban counterparts: a comparative, time-series (1997–2008) analysis. *J Rural Health* 2011;27:270–7.
- 99 Tokmakidis SP, Kasambalis A, Christodoulos AD. Fitness levels of Greek primary schoolchildren in relationship to overweight and obesity. *Eur J Pediatr* 2006;165:867–74.
- 100 Manios Y, Yiannakouris N, Papoutsakis C. Behavioral and physiological indices related to BMI in a cohort of primary schoolchildren in Greece. *Am J Hum Biol* 2004;16:639–47.
- 101 Georgiadis G. *Evaluation of physical fitness of Greek youth aged 6-18 years* [dissertation]. Athens, Greece: University of Athens, 1993.
- 102 Welk GJ, Saint-Maurice PF, Csányi T. Health-related physical fitness in Hungarian youth: age, sex, and regional profiles. *Res Q Exerc Sport* 2015;86:S45–57.
- 103 Barabás A. Measurement of aerobic power by field tests. In: Coudert J, Van Praagh E, eds. *Pediatric work physiology: children and exercise XVI*. Paris: Masson, 1992:39–41.
- 104 Gunnarsson HG, Sigríksson R. *Eurofit: physical fitness of Icelandic pupils at age of 6-15 years old*. Reykjavik, Iceland, 1999.
- 105 Grassi GP, Turci M, Sforza C. Aerobic fitness and somatic growth in adolescents: a cross sectional investigation in a high school context. *J Sports Med Phys Fitness* 2006;46:412–18.
- 106 Cilia G, Bazzano C, Bellucci M, *et al*. I risultati dei test Eurofit nella scuola Matteucci di Roma. *Alcmeone* 1998;2:16–20.
- 107 Cilia G, Bellucci M, Bazzano C, *et al*. Eurofit 1997: Banche dati per la scuola. *Alcmeone* 1997;3:13–32.
- 108 Cilia G, Bellucci M, Riva M. *Eurofit 1995*. Roma: Istituto Superiore Statale di Educazione Fisica, 1996.
- 109 Cilia G, Bellucci M. *Eurofit: tests Europei di attitudine fisica*. Roma: Istituto Superiore Statale di Educazione Fisica, 1993.
- 110 Council of Europe. *Évaluation de l'aptitude physique: Eurofit batterie expérimentale*. Rome: Council of Europe, 1986.
- 111 Ministry of Education, Culture, Sports, Science and Technology. *Report book on the survey of physical fitness and athletic ability*. Tokyo: Ministry of Education, Culture, Sports, Science and Technology, 1999–2015.

- 112 Sauka M, Priedite IS, Artjuhova L, *et al.* Physical fitness in northern European youth: reference values from the Latvian physical health in youth study. *Scand J Public Health* 2011;39:35–43.
- 113 Cazorla G, Dudal J, Garinet P. État de développement des capacités physiques des jeunes mauriciens 7-18 ans et plus. Rapport pour le Ministère Mauricien de la Jeunesse et Sports et pour le Ministère Français de la Coopération, 1986.
- 114 Galaviz KI, Tremblay MS, Colley R, *et al.* Associations between physical activity, cardiorespiratory fitness, and obesity in Mexican children. *Salud Pública Méx* 2012;54:463–9.
- 115 Dadouchi MF, Boudiab K, Yahia J, *et al.* *De la détermination du profil athlétique marocain au contenu et à l'évaluation.* Ministère de l'Éducation nationale et de la jeunesse. École normale supérieure de Marrakech, 2003.
- 116 Brouwer SI, Stol RP, Liem ET, *et al.* The role of fitness in the association between fatness and cardiometabolic risk from children to adolescence. *Pediatr Diabetes* 2013;14:57–65.
- 117 Slinger J, van Breda E, Kuipers H. Aerobic fitness data for Dutch adolescents (2002-2005). *Pediatr Exerc Sci* 2009;21:10–18.
- 118 van Mechelen W, van Lier WH, Hlobil H, *et al.* *Eurofit: Handleiding met referentieschalen voor 12-tot en met 16-jarige jongens en meisjes in Nederland.* Haarlem: Uitgeverij de Vrieseborch, 1991.
- 119 Haugen T, Høigaard R, Seiler S. Normative data of BMI and physical fitness in a Norwegian sample of early adolescents. *Scand J Public Health* 2014;42:67–73.
- 120 Cossio-Bolaños MA, Arruda M. Propuesta de valores normativos para la evaluación de la aptitud física en niños de 6 a 12 años de Arequipa, Perú. *Rev Med Herediana* 2009;20:206–12.
- 121 Gonzalez-Suarez CB, Grimmer-Sommers K. The association of physical activity and physical fitness with pre-adolescent obesity: an observational study in Metromanila, Phillipines. *J Phy Act Health* 2011;8:804–10.
- 122 Bronikowski M, Bronikowska M. Salutogenesis as a framework for improving health resources of adolescent boys. *Scand J Public Health* 2009;37:525–31.
- 123 Pilicz S, Przeweda R, Dobosz J, *et al.* *Punktacja sprawnosci fizycznej mlodziezy Polskiej wg miedzynarodowego testu sprawnosci fizycznej: Kryteria pomiaru wydolnosci organizmu testem Coopera.* Warszawa: Akademia Wychowania Fizycznego Józefa Pilsudskiego, 2003.
- 124 Maciaszek J, Osinski W. Poziom sprawnosci fizycznej u chlopców i dziewczat Poznanskich w wieku 10-14 lat. *Rocz Naukowe AWF Poznaniu* 2001;50:3–17.
- 125 Mleczko E, Ozimek M. *Rozwój somatyczny i motoryczny mlodziezy Krakowskiej miedzy 15 a 19 rokiem zycia z uwzglednieniem czynników srodowiskowych.* Kraków: Akademia Wychowania Fizycznego, 2000.
- 126 Santos R, Mota J, Santos DA, *et al.* Physical fitness percentiles for Portuguese children and adolescents aged 10-18 years. *J Sports Sci* 2014;32:1510–81.
- 127 Coelho-Silva MJ, Ronque ERV, Cyrino ES, *et al.* Nutritional status, biological maturation and cardiorespiratory fitness in Azorean youth aged 11-15 years. *BMC Public Health* 2013;13:495.

- 128 Rodrigues LP, Leitaó R, Lopes VP. Physical fitness predicts adiposity longitudinal changes over childhood and adolescence. *J Sci Med Sport* 2013;16:118–23.
- 129 Marta CC, Marinho DA, Barbosa TM. Physical fitness differences between prepubescent boys and girls. *J Strength Cond Res* 2012;26:1756–66.
- 130 Mota J, Guerra S, Leandro C. Association of maturation, sex, and body fat in cardiorespiratory fitness. *Am J Hum Biol* 2002;14:707–12.
- 131 Kim H-B, Stebbins CL, Chai J-H, *et al.* Taekwondo training and fitness in female adolescents. *J Sports Sci* 2011;29:133–8.
- 132 Cazorla G, Dudal J, Faye J. *État de développement des capacités physiques des jeunes sénégalais 7-18 ans et plus. Rapport pour le Ministère Sénégalais de la Jeunesse et Sports et pour le Ministère Français de la Coopération.* Paris, France: Ministère Sénégalais de la Jeunesse et Sports et pour le Ministère Français de la Coopération, 1988.
- 133 Bovet P, Auguste R, Burdette H. Strong inverse association between physical fitness and overweight in adolescents: a large school-based survey. *Int J Behav Nutr Phys Act* 2007;4:24.
- 134 Cazorla G, Rousseau G, Dudal J, *et al.* *Évaluation des capacités motrices de l'enfant, de l'adolescent et du jeune seychellois : 7-18 ans et plus. Rapport pour le Ministère Seychellois de l'Éducation et pour le Ministère Français de la Coopération.* Paris, France: Ministère Seychellois de l'Éducation et Ministère Français de la Coopération, 1990.
- 135 Krska P, Sedláček J, Hubinák A, *et al.* General motor fitness and somatic parameters comparison between former population and present primary school girls in Ruzomberok. In: Proceedings of the 9th International Conference in Physical Education, Sport and Physical Therapy, Iași, 13–14 November. Iași, Romania: Faculty of Sports and Physical Education, Alexandru Ioan Cuza University, 2015.
- 136 Krska P, Sedláček J, Hubinák A, *et al.* Physical development and general motor performance of present primary school boys population in Ruzomberok. In: Zvonář M, Sajdlová Z, eds. Proceedings of the 10th International Conference of Kinanthropology, Brno, Czech Republic, 18–20 November. Brno: Faculty of Sport Studies, Masaryk University, 2015:289–96.
- 137 Kyselovicová O. Programy aerobiku z aspektu rozvoja telesnej zdatnosti dievcat na gymnáziu. *Acta Facultatis Educationis Physicae Univ Comenianae* 2000;41:50–61.
- 138 Kasa J, Majherová M. Physical and motor development of children by Eurofit. *Stud Psychol (Bratisl)* 1997;39:270–4.
- 139 Moravec R. Eurofit—physique and motor fitness of the Slovak school youth. In: Moravec R, Kampmiller T, Sedláček J, eds. *Eurofit—physique and motor fitness of the Slovak school youth.* Bratislava: Slovak Scientific Society for Physical Education and Sports, 1996:9–22.
- 140 Belej M, Junger J, Mikus M. Vykonnosť žiakov prijatých na stredné školy vo východoslovenskom regióne zisťovaná systémom EUROFIT. *Tel Vych Sport* 1995;5:12–15.
- 141 Pienaar C, Coetzee B, Monyeki AM. The use of anthropometric measurements and the influence of demographic factors on the prediction of VO₂max in a cohort of adolescents: the PAHL study. *Ann Hum Biol* 2015;42:135–43.

- 142 Du Toit D, Pienaar AE, Truter L. Relationship between physical fitness and academic performance in South African children. *S Afr J Res Sport PH* 2011;33:23–35.
- 143 Pienaar AE, Viljoen A. Physical and motor ability, anthropometrical and growth characteristics of boys in the northwest province of South Africa: a sport talent perspective. *S Afr J Res Sport Phys Educ Recreation* 2010;32:71–93.
- 144 Du Preez SM. *The effect of physical activity on the body composition and health related fitness of 9 to 13-year-old boys* [thesis]. Potchefstroom, South Africa: North-West University, 2008.
- 145 Stadler MC. *The influence of a physical activity intervention program (PAI) on the physical fitness levels, body composition and health risk behaviour of 9 to 13 year old girls* [dissertation]. Potchefstroom, South Africa: North-West University, 2007.
- 146 Du Toit L, Venter RE, Potgieter JR. The relationship between cardiorespiratory fitness, body composition and physical self-perception of adolescent girls. *J Hum Mov Stud* 2005;48:353–64.
- 147 Van Gent M, Malan DDJ, Pienaar AE. A comparison of the anthropometric, physical and motor growth characteristics of 12-15 year old girls in the North West province with Australian girls. *Afr J Phys Health, Educ Recr Dance* 2002;8:309–20.
- 148 Du Randt R. *The 1996 South African sport talent identification project: report prepared for the Sports Information and Science Agency*. Pretoria: South African Sports Commission, 1996.
- 149 Gullías-González R, Martínez-Vizcaíno V, García-Prieto JC, *et al.* Excess of weight, but not underweight, is associated with poor physical fitness in children and adolescents from Castilla-La. *Eur J Pediatr* 2014;173:727–35.
- 150 Torrijos-Niño C, Martínez-Vizcaíno V, Pardo-Guijarro MJ, *et al.* Physical fitness, obesity, and academic achievement in schoolchildren. *J Pediatr* 2014;165:104–9.
- 151 Chillón P, Ortega FB, Ferrando JA, *et al.* Physical fitness in rural and urban children and adolescents from Spain. *J Sci Med Sport* 2011;14:417–23.
- 152 Castro-Pinero J, González-Montesinos JL, Mora J, *et al.* Percentile values for muscular strength field tests in children aged 6 to 17 years: influence of weight status. *J Strength Cond Res* 2009;23:2295–310.
- 153 Ortega FB, Ruiz JR, Castillo MJ, *et al.* Low level of physical fitness in Spanish adolescents. Relevance for future cardiovascular health (AVENA Study). *Rev Esp Cardiol* 2005;58:898–909.
- 154 García Baena J. *La condición física en la educación secundaria. Trabajo de investigación* [thesis]. Madrid, Spain: Universidad Nacional de Educacion a Distancia, 1999.
- 155 Prat JA, Casamort J, Balagué N, *et al.* Eurofit: *La batería Eurofit en Catalunya*. Barcelona: Secretaria General de l'Esport, 1998.
- 156 Tercedor P, Delgado-Fernandez M. Condicion física relacionada con la salud en escolares de 10 años de edad de Granada. *Proceedings of the II congreso internacional sobre la enseñanza de la educación física y el deporte escolar (second international congress about teaching physical education and school sport)*. 1998.

- 157 Sainz RM. *La batería Eurofit en Euskadi*. Vitoria-Gasteiz: Instituto Vasco de Educación Física, 1996.
- 158 Ureña F. *Valoración y baremación de la aptitud física en el alumnado de segundo ciclo de educación secundaria obligatoria de la comunidad autónoma de Murcia. Su utilización según los postulados de la reforma* [dissertation]. Murcia, Spain: Universidad de Murcia, 1996.
- 159 Brito Ojeda EM, Navarro Valdivielso M, García Afonso D, *et al.* *La condición física en la población escolar de gran Canaria (10-19 años)*. Las Palmas de Gran Canaria, Spain: Excmo. Cabildo Insular de Gran Canaria, 1995.
- 160 Sainz RM. *Aptitudes psíquicas y físicas: Estudio de la aptitud física de los adolescentes de la provincia de Vizcaya y su relación con la personalidad* [dissertation]. Bilbao, Spain: Universidad de Deusto, 1992.
- 161 Rivas FJ. Valoración de la evolución antropométrica y de las características motrices en la población escolar de un centro de E.G.B. mediante un estudio transversal. In: *II Congreso Galego da Educación Física e o Deporte*. La Coruña: Escola Galega do Deporte de la Xunta de Galicia, 1987:17–28.
- 162 Lieveld J, Vrijens J, Bouckaert J. Ethnic differences in body structure and physical fitness in Surinamese boys aged 14 years. In: Claessens AL, Lefevre J, Vanden Eynde B, eds. *World-wide variation in physical fitness*. Leuven: Katholieke Universiteit Leuven, 1993:87–92.
- 163 Schmid M, Romann M, Kriemler S, *et al.* Wie kann die Fitness von Schulkindern gemessen werden? *Sportmedizin Sporttraumatologie* 2007;55:52–61.
- 164 Caudey M, Narring F, Michaud P-A. A cross-sectional survey assessing physical fitness of 9- to 19-year-old girls and boys in Switzerland. *Pediatr Exerc Sci* 2000;12:398–412.
- 165 Aandstad A, Berntsen S, Hageberg R, *et al.* A comparison of estimated maximal oxygen uptake in 9 and 10 year old schoolchildren in Tanzania and Norway. *Br J Sports Med* 2006;40:278–92.
- 166 Çalis M, Ergen E, Turnagöl H, *et al.* Beden egitimi derslerinin bir öğretim yılı boyunca 15-16 yaş grubu öğrenciler üzerindeki fizyolojik etkilerinin Eurofit test bataryası ile izlenmesi. In: *3 Ulusal Spor Bilimleri Kongresi*. Ankara, Turkey: Hacettepe Üniversitesi, 1992:367–72.
- 167 Richards J, Foster C, Townsend N. Physical fitness and mental health impact of a sport-for-development intervention in a post-conflict setting: randomised controlled trial nested within an observational study of adolescents in Gulu, Uganda. *BMC Public Health* 2014;14:619.
- 168 Sandercock G, Ogunleye A, Voss C. Six-year changes in body mass index and cardiorespiratory fitness of English schoolchildren from an affluent area. *Int J Obes* 2015;39:1504–7.
- 169 Sandercock G, Voss C, Cohen D, *et al.* Centile curves and normative values for the twenty metre shuttle-run test in English schoolchildren. *J Sports Sci* 2012;30:679–87.
- 170 Boddy L, Fairclough SJ, Atkinson G, *et al.* Changes in cardiorespiratory fitness in 9- to 10.9-year-old children: Sports Linx 1998-2010. *Med Sci Sports Exerc* 2012;44:481–6.
- 171 Sandercock G, Voss C, McConnell D. Ten year secular declines in the cardiorespiratory fitness of affluent English children are largely independent of changes in body mass index. *Arch Dis Child* 2010;95:46–7.

- 172 Liverpool City Council. *Liverpool Sports Linx Project 01-03: Report on the health and fitness of Liverpool primary and secondary school children*. Liverpool, UK: Liverpool City Council, 2003.
- 173 Boreham C, Twisk J, Murray L, *et al*. Fitness, fatness, and coronary heart disease risk in adolescents: the Northern Ireland Young Hearts Project. *Med Sci Sports Exerc* 2001;33:270–4.
- 174 Watkins DC. *Ten year trends (1990-2000) in biological and behavioural risk factors for coronary heart disease in northern Irish adolescents* [thesis]. Belfast, UK: The Queen's University of Belfast, 2001.
- 175 Twisk JWR, Boreham C, Cran G, *et al*. Clustering of biological risk factors for cardiovascular disease and the longitudinal relationship with lifestyle of an adolescent population: the Northern Ireland Young Hearts Project. *J Cardiovasc Risk* 1999;6:355–62.
- 176 Mahoney C. 20-MST and PWC170 validity in non-Caucasian children in the UK. *Br J Sports Med* 1992;26:45–7.
- 177 Mahoney CA, Boreham CAG. Eurofit in Belfast primary schools. *Scot J Phys Educ* 1991;19:1–4.
- 178 Riddoch C, Savage JM, Murphy N, *et al*. Long term health implications of fitness and physical activity patterns. *Arch Dis Child* 1991;66:1426–33.
- 179 Boreham CAG, Paliczka VJ, Nichols AK. Fitness testing of Belfast schoolchildren. In: 5th European research seminar on testing physical fitness. *Strasbourg: Council of Europe*, 1987:52–7.
- 180 Lewitt MS, Baker JS, Mooney GP, *et al*. Pubertal stage and measures of adiposity in British schoolchildren. *Ann Hum Biol* 2012;39:440–7.
- 181 Ranson R, Stratton G, Taylor S. Digit ratio (2D:4D) and physical fitness (Eurofit test battery) in school children. *Early Hum Dev* 2015;91:327–31.
- 182 Liu W, Zillifro TD, Nichols RA. Tracking of health -related physical fitness for middle school boys and girls. *Pediatr Exerc Sci* 2012;24:549–62.
- 183 Welk GJ, Maduro PFDS-M, Laurson KR, *et al*. Field evaluation of the new FITNESSGRAM criterion-referenced standards. *Am J Prev Med* 2011;41:S131–42.
- 184 Beets MW, Pitetti KH, Cardinal BJ. Progressive aerobic cardiovascular endurance run and body mass index among an ethnically diverse sample of 10–15 year olds. *Res Q Exerc Sport* 2005;76:389–97.
- 185 Beets MW, Pitetti KH. A comparison of shuttle-run performance between mid-western youth and their national and international counterparts. *Pediatr Exerc Sci* 2004;16:94–112.
- 186 Welk GJ, Schaben JA, Shelley M. Physical activity and physical fitness in children schooled at home and children attending public schools. *Pediatr Exerc Sci* 2004;16:310–23.
- 187 Lloyd LK, Bishop PA, Walker JL, *et al*. The influence of body size and composition on FITNESSGRAM® test performance and the adjustment of FITNESSGRAM® test scores for skinfold thickness in youth. *Meas Phys Educ Exerc Sci* 2003;7:205–26.
- 188 Chun DM, Corbin CB, Pangrazi RP. Validation of criterion-referenced standards for the mile run and progressive aerobic cardiovascular endurance tests. *Res Q Exerc Sport* 2000;71:125–34.

- 189 Wolford N. *The difference in physical fitness levels of fifth graders according to socioeconomic groups and genders* [dissertation]. Ann Arbor, MI: University Microforms International, 1998.
- 190 Mahar MT, Rowe DA, Parker CR, et al. Criterion-referenced and norm-referenced agreement between the mile run/walk and PACER. *Meas Phys Educ Exerc Sci* 1997;1:245–58.
- 191 <http://data.worldbank.org/about/country-and-lending-groups> (accessed 24 Dec 2015).
- 192 Cole TJ, Freeman JV, Preece MA. Body mass index reference curves for the UK, 1990. *Arch Dis Child* 1995;73:25–9.
- 193 Cole TJ, Bellizzi MC, Flegal KM, et al. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* 2000;320:1240–3.
- 194 Eisenmann JC. Waist circumference percentiles for 7- to 15-year-old Australian children. *Acta Paediatr* 2005;94:1182–5.
- 195 Fernandez JR, Redden DT, Pietrobelli A, et al. Waist circumference percentiles in nationally representative samples of African-American, European-American, and Mexican-American children and adolescents. *J Paediatr* 2004;145:439–44.
- 196 McCarthy HD, Jarret KV, Crawley HF. The development of waist circumference percentiles in British children aged 5.0–16.9 y. *Eur J Clin Nutr* 2001;55:902–7.
- 197 Shields M. Overweight and obesity among children and youth. *Health Rep* 2006;17:27–42.
- 198 Tremblay MS, Shields M, Laviolette M, et al. Fitness of Canadian children and youth: results from the 2007–2009 Canadian Health Measures Survey. *Health Rep* 2009;21:1–14.
- 199 Jackson LV, Thalange NKS, Cole TJ. Blood pressure centiles for Great Britain. *Arch Dis Child* 2007;92:298–303.
- 200 Jolliffe CJ, Janssen I. Development of age-specific adolescent metabolic syndrome criteria that are linked to the Adult Treatment Panel III and International Diabetes Federation criteria. *J Am Coll Cardiol* 2007;49:891–8.
- 201 Castro-Piñero J, González-Montesinos JL, Keating XD, et al. Percentile values for running sprint field tests in children ages 6–17 years: influence of weight status. *Res Q Exerc Sport* 2010;81:143–51.
- 202 De Miguel-Etayo P, Gracie-Marco L, Ortega FB, et al. Physical fitness reference standards in European children: the IDEFICS study. *Int J Obes* 2014;38 2):S57–66.
- 203 Eisenmann JC, Laurson KR, Welk G J. Aerobic fitness percentiles for U.S. adolescents. *Am J Prev Med* 2011;41:S106–10.
- 204 Armstrong N, Barker AR. Endurance training and elite young athletes. *Med Sport Sci* 2011;56:59–83.
- 205 Ruiz JR, Ortega FB, Rizzo NS, et al. High cardiovascular fitness is associated with low metabolic risk score in children: the European Youth Heart Study. *Pediatr Res* 2007;61:350–5.
- 206 Lobelo F, Pate RR, Dowda M, et al. Validity of cardiorespiratory fitness criterion-referenced standards for adolescents. *Med Sci Sports Exerc* 2009;41:1222–9.

- 207 Adegboye ARA, Anderssen SA, Froberg K, *et al.* Recommended aerobic fitness level for metabolic health in children and adolescents: a study of diagnostic accuracy. *Br J Sports Med* 2011;45:722–8.
- 208 Boddy LM, Thomas NE, Fairclough SJ, *et al.* ROC generated thresholds for field-assessed aerobic fitness related to body size and cardiometabolic risk in schoolchildren. *PLoS ONE* 2012;7:e45755.
- 209 Mesa JL, Ruiz JR, Ortega FB, *et al.* Aerobic fitness in relation to blood lipids and fasting glycaemia in adolescents: influence of weight status. *Nutr Metab Cardiovasc Dis* 2006;16:285–93.
- 210 Moreira C, Santos R, Ruiz JR, *et al.* Comparison of different VO₂max equations in the ability to discriminate the metabolic risk in Portuguese adolescents. *J Sci Med Sport* 2011;14:79–84.
- 211 Ruiz JR, Huybrechts I, Cuenca-Garcia M, *et al.* Cardiorespiratory fitness and ideal cardiovascular health in European adolescents. *Heart* 2015;101:766–73.
- 212 Tomkinson G. Aerobic fitness thresholds for cardiometabolic health in children and adolescents. *Br J Sports Med* 2011;45:686–7.
- 213 Barnett A, Chan LYS, Bruce IC. A preliminary study of the 20-m multistage shuttle run as a predictor of peak VO₂ in Hong Kong Chinese students. *Pediatr Exerc Sci* 1993;5:42–50.
- 214 Stickland MK, Petersen SR, Bouffard M. Prediction of maximal aerobic power from the 20-m multistage shuttle run test. *Can J Appl Physiol* 2003;28: 272–82.
- 215 Flouris AD, Metsios GS, Koutedakis Y. Enhancing the efficacy of the 20 m multistage shuttle run test. *Br J Sports Med* 2005;39:166–70.
- 216 Matsuzaka A, Takahashi Y, Yamazoe M, *et al.* Validity of the multistage 20-m shuttle-run test for Japanese children, adolescents, and adults. *Pediatr Exerc Sci* 2004;16:113–25.
- 217 Mahar MT, Welk GJ, Rowe DA, *et al.* Development and validation of a regression model to estimate VO₂peak from PACER 20-m shuttle run performance. *J Phys Act Health* 2006;3:S34–46.
- 218 Ruiz JR, Ramirez-Lechuga J, Ortega FB, *et al.* Artificial neural network-based equation for estimating VO₂max from the 20 m shuttle run test in adolescents. *Artif Intell Med* 2008;44:233–45.
- 219 Oliveira J. *Validação directa do teste de vaivém em 20 metros, de Luc- Léger, em adolescentes portuguesas*. Cruz-Quebrada: FMH-UTL, 1998.
- 220 Pitetti KH, Fernhall B, Figoni S. Comparing two regression formulas that predict VO₂peak using the 20-m shuttle run for children and adolescents. *Pediatr Exerc Sci* 2002;14:125–34.
- 221 McVeigh SK, Payne AC, Scott S. The reliability and validity of the 20-meter shuttle test as a predictor of peak oxygen uptake in Edinburgh school children, age 13 to 14 years. *Pediatr Exerc Sci* 1995;7:69–79.
- 222 Moreno LA, González-Gross M, Kersting M, *et al.* Assessing, understanding and modifying nutritional status, eating habits and physical activity in European adolescents: the HELENA (Healthy lifestyle in Europe by nutrition in adolescence) study. *Public Health Nutr* 2008;11:288–99.

- 223 Andersen KL, Seliger V, Rutenfranz J, *et al.* Physical performance capacity of children in Norway. Part IV. The rate of growth in maximal aerobic power and the influence of improved physical education of children in a rural community—population parameters in a rural community. *Eur J Appl Physiol Occup Physiol* 1976;35:49–58.
- 224 Armstrong N, Welsman JR. Assessment and interpretation of aerobic fitness in children and adolescents. *Exerc Sport Sci Rev* 1994;22:435–76.
- 225 Krahenbuhl GS, Skinner JS, Kohrt WM. Developmental aspects of maximal aerobic power in children. *Exerc Sport Sci Rev* 1985;13:503–38.
- 226 Rowland TW. Evolution of maximal oxygen uptake in children. *Med Sport Sci* 2007;50:200–9.
- 227 Rowland TW. *Children's exercise physiology*. 2nd edn. Champaign, IL: Human Kinetics, 2005:89–112.
- 228 Bar-Or O, Rowland TW. Physiologic and perceptual responses to exercise in the healthy child. In: Bar-Or O, Rowland TW, eds. *Pediatric exercise medicine: from physiologic principles to health care application*. Champaign, IL: Human Kinetics, 2004:3–59.
- 229 Tomkinson G, Macfarlane DJ, Noi S, *et al.* Temporal changes in long-distance running performance of Asian children between 1964 and 2009. *Sports Med* 2012;42:267–79.
- 230 Moliner-Urdiales D, Ruiz JR, Ortega FB, *et al.* Secular trends in health-related physical fitness in Spanish adolescents: the AVENA and HELENA Studies. *J Sci Med Sport* 2010;13:584–8.
- 231 Tomkinson G, Olds T, Borms J. Who are the Eurofittest? *Med Sport Sci* 2007;50:104–28.
- 232 Castro-Piñero J, Artero EG, España-Romero V, *et al.* Criterion-related validity of field-based fitness tests in youth: a systematic review. *Br J Sports Med* 2010;44:934–43.
- 233 Mayorga-Vega D, Aguilar-Soto P, Viciano J. Criterion-related validity of the 20-M Shuttle run test for estimating cardiorespiratory fitness: a meta-analysis. *J Sports Sci Med* 2015;14:536–47.
- 234 Lussier L, Buskirk ER. Effects of an endurance training regimen of assessment of work capacity in prepubertal children. *Ann N Y Acad Sci* 1977;301:734–47.
- 235 Krahenbuhl GS, Pangrazi RP, Chomokos EA. Aerobic responses of young boys to submaximal running. *Res Q* 1979;50:413–21.
- 236 Péronnet F, Thibault G. Mathematical analysis of running performance and world records. *J Appl Physiol* 1989;67:453–65.
- 237 Sjödin B. The relationship among running economy, aerobic power, and onset of blood lactate accumulation in young boys (11–15 years). In: Komi PV, ed. *Exercise and sport biology*. Champaign, IL: Human Kinetics, 1982:57–60.
- 238 Mayers N, Gutin B. Physiological characteristics of elite prepubertal cross-country runners. *Med Sci Sports* 1979;11:172–6.
- 239 Cureton KJ, Boileau RA, Lohman TG, *et al.* Determinants of distance running performances in children: analysis of a path model. *Res Q* 1977;42:270–9.

- 240 Voss C, Sandercock G. Does the twenty meter shuttle-run test elicit maximal effort in 11- to 16-year-olds? *Pediatr Exerc Sci* 2009;21:55–62.
- 241 Longmuir PE, Colley RC, Wherley VA, *et al.* Canadian Society for Exercise Physiology position stand: benefit and risk for promoting childhood physical activity. *Appl Physiol Nutr Metab* 2014;39:1271–9.
- 242 Meusel D, Ruiz JR, Ortega FB, *et al.* Assessing levels of physical activity in the European population—the ALPHA project. *Seleccion* 2007;16:9–12.
- 243 Longmuir PE, Boyer C, Lloyd M, *et al.* The Canadian Assessment of Physical Literacy: methods for children in grades 4 to 6 (8 to 12 years). *BMC Public Health* 2015;15:767.
- 244 Council of Europe. *Eurofit: handbook for the EUROFIT tests of physical fitness*. Rome: Secretariat of the Committee for the Development of Sport within the Council of Europe, 1998.
- 245 Welk GJ, Going SB, Morrow Jr JR, *et al.* Development of new criterion-references fitness standards in the FITNESSGRAM® program: rationale and conceptual overview. *Am J Prev Med* 2011;41:S63–7.
- 246 Ortega FB, Cadenas-Sánchez C, Sánchez-Delgado G, *et al.* Systematic review and proposal of a field-based physical fitness-test battery in preschool children: the PREFIT battery. *Sports Med* 2015;45:533–55.
- 247 Cadenas-Sánchez C, Alcántara-Moral F, Sánchez-Delgado G, *et al.* [Assessment of cardiorespiratory fitness in preschool children: adaptation of the 20 metre shuttle run test]. *Nutr Hosp* 2014;30:1333–43.