



11-3-2018

Temporal trends in the cardiorespiratory fitness of 2,525,827 adults between 1967 and 2016

Nicholas R. Lamoureux

John S. Fitzgerald
University of North Dakota, john.s.fitzgerald@UND.edu

Kevin I. Norton

Todd Sabato
University of North Dakota, todd.sabato@und.edu

Mark S. Tremblay

See next page for additional authors

[How does access to this work benefit you? Let us know!](#)

Follow this and additional works at: <https://commons.und.edu/ehb-fac>



Part of the [Arts and Humanities Commons](#), and the [Health and Physical Education Commons](#)

Recommended Citation

Nicholas R. Lamoureux, John S. Fitzgerald, Kevin I. Norton, et al.. "Temporal trends in the cardiorespiratory fitness of 2,525,827 adults between 1967 and 2016" (2018). *Education, Health & Behavior Studies Faculty Publications*. 35.

<https://commons.und.edu/ehb-fac/35>

This Article is brought to you for free and open access by the Department of Education, Health & Behavior Studies at UND Scholarly Commons. It has been accepted for inclusion in Education, Health & Behavior Studies Faculty Publications by an authorized administrator of UND Scholarly Commons. For more information, please contact und.common@library.und.edu.

Authors

Nicholas R. Lamoureux, John S. Fitzgerald, Kevin I. Norton, Todd Sabato, Mark S. Tremblay, and Grant R. Tomkinson

Temporal Trends in the Cardiorespiratory Fitness of 2,525,827 Adults Between 1967 and 2016: A Systematic Review

Nicholas R. Lamoureux, John S. Fitzgerald, Kevin I. Norton, Todd Sabato, Mark S. Tremblay, Grant R. Tomkinson

Abstract

Objective To estimate international and national temporal trends in the cardiorespiratory fitness (CRF) of adults, and to examine relationships between trends in CRF and trends in health-related, socioeconomic, and environmental indicators.

Methods Data were obtained from a systematic search of studies that explicitly reported temporal trends in the CRF of apparently healthy adults aged 18–59 years. Sample-weighted temporal trends were estimated using best-fitting regression models relating the year of testing to mean CRF. Post-stratified population-weighted mean changes in percent and standardized CRF were estimated. Pearson's correlations were used to describe associations between linear trends in CRF and linear trends in health-related, socioeconomic, and environmental indicators.

Results 2,525,827 adults representing eight high- and upper-middle-income countries between 1967 and 2016 collectively showed a moderate decline of 7.7% (95% CI –8.4 to –7.0) or 1.6% per decade (95% CI –1.7 to –1.5). Internationally, CRF improved in the 1960s and 1970s, and progressively declined at an increasing rate thereafter. Declines were larger for men than for women, and for young adults (<40 years) than for middle-aged adults (≥ 40 years). All countries experienced declines in CRF with a very strong negative correlation between CRF trends and obesity trends.

Conclusions There has been a meaningful decline in the CRF of adults since 1980, which has progressively increased in magnitude over time, suggestive of a corresponding decline in population health. Continuous national and international surveillance systems are needed in order to monitor health and fitness trends, especially among low- and middle-income countries for which data do not currently exist. PROSPERO registration number: CRD42013003678.

1 Introduction

Cardiorespiratory fitness (CRF) is an important marker of cardiovascular health [1, 2] as it provides a measure of the body's capacity to deliver and utilize oxygen for energy transfer to support muscle activity during physical activity and exercise [3]. CRF—the fourth-leading risk factor for cardiovascular disease [4]—is strongly and inversely related (independent of physical activity level and weight status) in adults to all-cause and cardiovascular disease mortality [5–12], stroke [9], diabetes [13], several cancers [14–17], and numerous other cardiovascular disease risk factors [18–20]. CRF has been proposed as an important clinical vital sign because of the improved cardiovascular disease mortality risk classification (e.g., net reclassification improvement [NRI], and Framingham and European risk scores) when added to traditional biomarkers [1, 21–24]. Attributable fraction estimates from the Aerobics Center Longitudinal Study of 53,785 men and women showed that CRF was more strongly associated with all-cause mortality than traditional risk factors such as obesity, smoking, hypertension, high cholesterol, and diabetes [25]. A recent scientific statement from the American Heart Association (AHA) reported that every metabolic equivalent (MET; 3.5 ml/kg/min) increase in adult CRF was associated with a 10–25% decrease in age-adjusted all-cause and/or cardiovascular disease mortality [1]. This evidence provides good support for

the monitoring of temporal trends in CRF, which shed meaningful light on temporal trends in population health.

CRF is often expressed as maximal oxygen uptake ($\dot{V}O_{2max}$) or METs, and can be measured using a variety of maximal or sub-maximal field- or laboratory-based running, cycling, or stepping tests. Although laboratory tests of CRF (e.g., $\dot{V}O_{2max}$ measured using expired gas-analysis indirect calorimetry) directly measure the function of the physiological systems (cardiovascular, respiratory, metabolic, and neuromuscular) to perform prolonged continuous, rhythmic, dynamic, large muscle group physical activity of moderate to high intensity, they are impractical for population-based testing. On the other hand, field tests of CRF that provide a single measure that assesses the integrated responses of the physiological systems required to perform prolonged exercise in conventional physical activity conditions, offer a simple, feasible, practical, reliable, and valid alternative [26]. Much of what is known about temporal trends in CRF comes from children and adolescents, where schools have provided opportunities for population-based testing that do not typically exist for adults. Collectively, several large systematic reviews on temporal trends in the CRF of children and adolescents have shown an international decline from the mid-1970s to the early 2000s, followed by a plateau [27–29]. Data on 129,882 children and adolescents aged 6–19 years from 11 countries showed a decline in mean CRF of 4.3% per decade between 1981 and 2000 [27], with a follow-up study showing a decline of 3.6% per decade in 25.4 million 6- to 19-year-olds from 27 countries between 1958 and 2003 [28]. More recently, Tomkinson et al. [29] reported that the international decline in CRF has diminished and stabilized with negligible change since 2000 in 965,264 children aged 9–17 years from 19 high- and upper-middle-income countries. Unfortunately, little is known about current national and international trends in the CRF of adults. Furthermore, our recent systematic review identified a strong negative association between country-level temporal trends in children’s CRF and temporal trends in income inequality (Gini index) [29], meaning countries with a widening gap between rich and poor residents had less favorable trends (i.e., large declines) in CRF. Examining the associations between trends in adult CRF and trends in health-related, socioeconomic, and environmental indicators could provide further insight into the importance of these indicators and their potential population health implications.

This study extends our published research on temporal trends in the CRF of children and adolescents [27–29] to adults. The primary aim therefore was to systematically analyze national and international temporal trends in the CRF of adults. The secondary aim was to examine relationships between temporal trends in CRF and temporal trends in health-related, socioeconomic, and environmental indicators across countries. We hypothesized that adult CRF has declined in recent decades, and that country-level temporal trends in health-related/socioeconomic/environmental indicators would be meaningfully associated with temporal trends in CRF.

2 Methods

2.1 Protocol and Registration

The review protocol was prospectively registered with the International Prospective Register of Systematic Review (PROSPERO; registration number CRD42013003678). This review was conducted and reported in accordance with the Preferred Reporting Items for Systematic review and Meta-analysis Protocols (PRISMA-P) statement for reporting systematic reviews [30].

2.2 Eligibility Criteria

Studies were included if they reported on temporal trends in the CRF of apparently healthy (free from known disease/injury) adults (aged 18 years or older) across at least two time points spanning a minimum

of 10 years. All studies must have relied on population-representative national fitness surveys of adults, with a minimum sample size of 100 per country-sex-age-test group. CRF must have been measured using non-gas-analyzed laboratory tests (e.g., submaximal bench stepping or cycling tests) or field tests (e.g., distance running/walking tests), with results reported as trends in estimated maximum oxygen uptake (estimated from submaximal bench stepping or cycling) or long-distance running performance (e.g., the time taken to run/walk a long distance or the distance run/walked over a long time).

2.3 Information Sources

A systematic literature search was conducted on 1 March 2018 using the EBSCO interface in Cumulative Nursing and Allied Health Literature (CINAHL), Educational Resources Information Center (ERIC), MEDLINE, and SPORTDiscus without date or language restrictions. The search strategy was developed with the assistance of an academic librarian experienced in systematic literature searching.

2.4 Search

The search was performed limited to abstract, title, and keywords. Search terms within a group were combined with the Boolean OR and were searched in conjunction with other search groups connected by the Boolean AND. Some terms were searched using proximity operators to search for the root word. The first group of search terms identified the fitness measure (physical fitness, or muscular strength, or muscular endurance, or aerobic fitness, or cardio* fitness or cardio* endurance). The second group identified the population (adult*, or men, or man, or woman, or women, or male or female). The third group identified the trend over time (secular, or temporal, or historical). Only studies published in English were included. The full search strategies for each database are shown in Electronic Supplementary Material 1.

2.5 Study Selection

Two researchers searched all databases independently, with all bibliographic records imported into RefWorks (version 2.0; ProQuest LLC, Ann Arbor, MI, USA) and de-duplicated. Both researchers independently screened all potentially relevant titles and abstracts against inclusion criteria, with exclusion by both researchers required for final exclusion. Full text copies were obtained and independently screened by two researchers against inclusion criteria, with consensus required for final inclusion. A third researcher resolved discrepancies if consensus was not reached. The reference lists of all included studies and personal libraries of the authors were reviewed to identify relevant studies not identified through the database search. In addition, while systematic reviews were not included in the analysis, their reference lists were reviewed for relevant studies. Where necessary, email contact was made with the corresponding authors in order to clarify published results, to avoid “double counting” previously reported data, or for additional information.

2.6 Data Collection Process

Descriptive data were extracted into a spreadsheet by a single researcher using a standardized study-specific template [29], and checked for accuracy by a second reviewer.

2.7 Data Items

The following study-specific descriptive data were extracted: title, country, years of testing, sex, age or age range, as well as the sample size, mean, and standard deviation data for measured CRF or the absolute, percent, and/or standardized changes in mean CRF ($\pm 95\%$ confidence intervals [CIs]). $\dot{V}O_{2\max}$ data were expressed as relative $\dot{V}O_{2\max}$ in ml/kg/ min, and long-distance running data were expressed as

running speeds in m/s because relative $\dot{V}O_2$ and $\dot{V}O_{2max}$ varies linearly with speed and maximal speed, and therefore speed should appropriately reflect $\dot{V}O_2$ (i.e., the underlying oxygen cost required to complete the distance run) [26, 31].

2.8 Summary Measures and Synthesis of Results

Temporal trends in mean CRF were analyzed using the detailed procedure described elsewhere [28, 29, 32]. Temporal trends at the country-sex-age-test level (e.g., 20- to 24-year-old Japanese women tested on the 1000-m run) were analyzed using best-fitting sample-weighted linear or polynomial (quadratic or cubic) regression models relating the year of testing to CRF. Changes in mean CRF were expressed as percent (% per year) changes. To interpret the magnitude of change, standardized effect sizes (ES) of 0.2, 0.5, and 0.8 were used as thresholds for small, moderate, and large, respectively, with $ES < 0.2$ considered to be negligible and $ES \geq 0.2$ considered to be meaningful. Positive changes indicated increases in mean CRF and negative changes indicated declines.

The temporal trends were described as follows: starting with the first year (Y_1) covered by any study-country-sex-age group, every group including Y_1 in its span of years was located, with every change (dx_1 , e.g., ml/kg/min or % per year) recorded [28, 29, 32]. This process was applied to all years for which change data were available ($Y_1 \dots Y_n$), yielding a series of yearly changes. The post-stratified population-weighted mean yearly change was calculated for year (Y_1) [33], which was repeated for $Y_2, Y_3, Y_4 \dots$ until the last year covered by any study, Y_n . This process yielded a series of population-weighted mean yearly changes (dx_1, \dots, dx_n) that collectively described the temporal pattern of change; a process that was performed for all adults and for different sex, age, and country groups. Population estimates were standardized to the year 1990—a common testing year across all country-sex-age groups—using United Nations data [34]. The post-stratification population-weighting procedure helped to correct the trends for systematic bias associated with over- and under-sampling, and to standardize the trends to underlying country-sex-age-specific demographics. Temporal trends were estimated using best-fitting population-weighted linear or polynomial regression models relating the year of testing to the yearly changes, and were graphically illustrated using an iterative procedure described by Tomkinson and Olds [28].

Relationships between linear temporal trends in CRF and linear temporal trends in broad health-related, socioeconomic, and environmental indicators across countries were quantified using Pearson's correlation coefficients, with 95% CIs estimated using Fisher's z-transformation. National trends for four health-related (prevalence of adult obesity [35]), socioeconomic (Human Development Index [HDI] [36]), and environmental (air pollution and urbanization [37]) indicators were analyzed using linear regression models (as described above for CRF). While several other health-related, socioeconomic, and environmental indicators could have been potentially explored (e.g., Gini index [a population measure of the distribution of wealth], physical activity levels), only trends in aforementioned indicators were selected because it was possible to calculate temporal trends using the same criteria as for CRF (e.g., across at least two time points spanning a minimum of 10 years) across the majority of included countries. To interpret the magnitude of correlation, ES of 0.1, 0.3, 0.5, 0.7, and 0.9 were used as thresholds for weak, moderate, strong, very strong, and nearly perfect, respectively, with $ES < 0.1$ considered to be negligible and $ES \geq 0.1$ considered to be meaningful.

3 Results

Figure 1 outlines the identification of the included studies. A total of 578 unique records were identified through online bibliographic database searching. After screening titles and abstracts, 19 articles were

retained for full-text review, of which 14 were excluded as described in Fig. 1. Five references plus three large population-representative national datasets (obtained from the personal library of the senior author) suitable for temporal trends analysis were retained for analysis.




Fig. 1 PRISMA flow chart outlining the flow of studies through the review

Temporal trends in CRF were estimated from 2,525,827 adults aged 18–59 years from eight countries (1113 country-sex-age-year groups) between 1967 and 2016 (Tables 1 and 2). These countries represented seven very high human development (high-income) countries and one high human development (upper-middle-income) country [36, 37], from three continents and approximately 23% of the world’s population. Trends were calculated for 60 country-sex-age groups (men: 31; women: 29; young adults [<40 years]: 46; middle-aged adults [≥ 40 years]: 14), with an average sample size of 42,097 (range 149–387,088) across an average span of 29 years (range 10–47).

Collectively, there was a moderate decline in mean CRF between 1967 and 2016 (change in means [95% CI]: -7.7% [-8.4 to -7.0]; ES -0.66 [-0.71 to -0.60]) (Fig. 2). There was a large collective decline in mean CRF in men (change in means [95% CI]: -10.0% [-10.9 to -9.1]; ES -0.83 [-0.90 to -0.77]), small declines in women (change in means [95% CI]: -5.3% [-6.2 to -4.4]; ES -0.48 [-0.56 to -0.40]) and young adults (change in means [95% CI]: -4.7% [-5.7 to -3.7]; ES -0.34 [-0.41 to -0.26]), and a negligible decline in middle-aged adults (change in means [95% CI]: -2.0% [-3.0 to -1.0]; ES -0.15 [-0.22 to -0.08]) (Fig. 3).

Internationally, the changes in mean CRF was not uniform over time, with the rate of change shifting from negligible improvements in the 1960s and 1970s (change in means [95% CI]: 0.4% per decade [0.2 – 0.6]; ES 0.05 [0.03 – 0.07]), to negligible declines in the 1980s and 1990s (change in means [95% CI]: -1.8% per decade [-2.0 to -1.6]; ES -0.16 [-0.17 to -0.15]), and to small declines in the 2000s and 2010s (change in means [95% CI]: -2.5% per decade [-2.8 to -2.2]; ES -0.23 [-0.25 to -0.21]) (Fig. 2). This trend was consistent across different sex and age groups (Fig. 3).

All countries experienced declines in CRF, ranging from large declines in China and Finland to negligible declines in Poland and the Republic of Korea (Fig. 2). While most country trends were approximately uniform (linear), Fig. 2 indicates non-uniform (curvilinear) trends in Finland and Poland evidenced by improvements in the 1970s and early 1980s followed by declines from the mid- to late-1980s onwards.

Table 1 Summary of the included studies by country

| Country | Sample information | Sex | Age (years) | Span of years | Sample size | Country-sex-age groups | HDI | CRF test(s) | Criterion validity (r) |
|-------------------------------------|---|------------------------|-------------|---------------|-------------|------------------------|-------------------|---|------------------------|
| Canada ^{w1} | Two national (stratified, random) surveys | M (48.1%) F (51.9%) | 20–59 | 1981–2009 | 9596 | 4 | 0.920 (very high) | Bench stepping (CAFT) ^a | 0.88 [43] |
| China ^{w2–8} | Seven national (stratified, random) surveys | M (50.7%) F (49.3%) | 18–22 | 1985–2014 | 499,229 | 10 | 0.738 (high) | Distance run (800–1000 m) ^b | 0.55–0.74 [42]* |
| Finland ^{w9} | 30 national (census) surveys | M | 20 | 1975–2004 | 387,088 | 1 | 0.895 (very high) | Timed run (12 min) ^c | 0.78 [42]* |
| Japan ^{w10–59} | 50 national (stratified, random) surveys | M (51.6%) F (48.4%) | 18–59 | 1967–2016 | 1,391,742 | 24 | 0.903 (very high) | Distance run/walk (1000–1500 m) ^b | 0.62–0.74 [42]*,† |
| Norway ^{w60} | Seven national (stratified, random) surveys | M | 18 | 1980–2002 | 184,638 | 1 | 0.949 (very high) | Stationary cycling (Åstrand-Rhyming) ^d | 0.83 [41] |
| Poland ^{w61} | Three national (stratified, random) surveys | M (53.1%) F (46.9%) | 18–19 | 1979–1999 | 33,358 | 4 | 0.855 (very high) | Distance run (800–1000 m) ^b | 0.55–0.74 [42]* |
| Republic of Korea ^{w62–69} | Eight national (stratified, random) surveys | M (56.2%) F (43.8%) | 19–49 | 1988–2007 | 19,207 | 12 | 0.901 (very high) | Distance run (1200 m) ^b | 0.74 [42]*,‡ |
| Singapore ^{w70} | Two national (stratified, random) surveys | M (42.3%) F (57.7%) | 18–19 | 1981–1991 | 969 | 4 | 0.925 (very high) | Distance run (2400 m) ^b | 0.79 [42]* |

CRF cardiorespiratory fitness, M male, F female, HDI Human Development Index (2015 estimate [36]) with HDI values of 0.800 and 0.700 used as thresholds for very high and high human development, respectively, CAFT Canadian Aerobic Fitness Test

a A multistage submaximal CRF stepping test performed on a double 20.3-cm step requiring participants to complete the necessary number of stages to reach a target heart rate of 85% of their age-predicted heart rate maximum

b A field-based running/walking test requiring participants to run/walk at the fastest pace possible without stopping around a fat marked outdoor course 800 m (China and Poland, women), 1000 m (China and Poland, men; Japan, women), 1200 m (Korea, men and women), 1500 m (Japan, men), or 2400 m (Singapore, men and women) in length

c A field-based running test requiring participants to run as far as possible without stopping around a fat marked outdoor course for 12 min

d A submaximal stationary cycling test requiring participants to pedal at an appropriately selected constant work rate for 6 min to determine steady-state heart rate; Criterion validity data are reported as correlation coefficients relating the CRF performance scores only to $\dot{V}O_{2\max}$ (ml/kg/min) measured using expired gas-analysis indirect calorimetry

*Overall weighted mean of r from a meta-analysis [42] where r was corrected for sampling error and measurement error

† Due to an absence of data, validity data for the 1600-m run/walk were assumed for the 1500-m run/walk

‡ Due to an absence of data, validity data for the 1000-m run/walk were assumed for the 1200-m run/walk

Country-level temporal trends for men and women were reasonably consistent, with women almost always experiencing smaller declines in CRF than men (e.g., Canada, China, Japan, and Poland) or improvements instead of declines (e.g., Republic of Korea) (Fig. 3). Similarly, country-level declines in CRF were consistently larger in young adults than in middle-aged adults (Fig. 3). There was a very strong negative correlation between trends in CRF and trends in adult obesity (r [95% CI] -0.77 [-0.96 to -0.03]), indicating that countries with the largest increases in obesity over the period 1990–2013 had the

largest declines in CRF (Table 3). The correlation between trends in CRF and trends in urbanization was strong and negative, and the correlations between trends in CRF and both trends in HDI and trends in air pollution were weak and positive (Table 3).




Fig. 2 National temporal trends in mean cardiorespiratory fitness (CRF) between 1967 and 2016. Data are standardized to the year 1990=100%, with higher values (>100%) indicating better CRF and negative values (<100%) indicating poorer CRF. The solid lines represent the national changes in mean CRF, with upward sloping lines indicating increases over time and downward sloping lines indicating declines over time. Mean (95% CI) percent changes (per decade) are shown at the top of each panel

4 Discussion

This analysis estimated that the CRF of adults: (a) improved internationally in the 1960s and 1970s and declined thereafter, with the magnitude of decline progressively increasing until it reached a peak in the 2000s and 2010s; (b) declined across all included countries, with both the rate and pattern of change varying among countries; and (c) declined to a greater extent in men than in women, and in younger adults than in middle-aged adults. It also indicated that countries with the largest increases in adult obesity experienced the largest declines in CRF. Given the strong and independent link between CRF and cardiovascular disease and all-cause mortality, these declines in CRF raise concern and suggest that the current and future health of today's adults is probably worse than that of their peers from decades past. For example, assuming a mean $\dot{V}O_{2max}$ of 45 ml/kg/min for young men [38, 39] and a 7.7% decline over the period 1967–2016, the reported international decline is equal to ~1 MET decline, which is associated with up to a 25% decrease in age-adjusted all-cause and/or cardiovascular disease mortality [1].

4.1 Temporal Declines in CRF

We have previously argued that temporal declines in CRF are influenced by a network of environmental, socioeconomic, behavioral, physical, psychosocial, and physiological variables [27–29]. Theoretically, declines in physiological variables such as $\dot{V}O_{2max}$ (leading to a reduced maximum rate at which oxygen can be delivered and utilized in energy transfer), mechanical efficiency (leading to an increased oxygen cost for any given exercise intensity), and the fractional utilization of oxygen (leading to a reduction in the length of time sustained for any given exercise intensity), as well as declines in psychosocial variables (e.g., motivation, ability to tolerate discomfort and pacing [with respect to long-distance running]), are

likely involved. While no population-representative trend data for these variables are available, analysis of 2006 Swedish adults aged 20–65 years between 1990 and 2000 showed that median absolute CRF ($\dot{V}O_{2max}$ in L/min estimated from the heart rate response to submaximal work on a stationary cycled) remained stable while median relative CRF ($\dot{V}O_{2max}$ in ml/kg/min) declined [40]. These data suggest that little change in true cardiovascular function (i.e., absolute $\dot{V}O_{2max}$) has occurred in adults over time, and that increased body mass—the ratio scaling factor used to remove the influence of body size on absolute $\dot{V}O_{2max}$ —is largely responsible for the decline in relative CRF. However, despite the fact that expired gas-analyzed measures of CRF were not used to track trends in CRF in this study, the included non-gas-analyzed measures demonstrate moderate-to-very high criterion validity against expired gas-analyzed $\dot{V}O_{2max}$, with validity coefficients ranging from 0.55 to 0.88 (Table 1). This suggests that the reported declines in CRF reflect declines in underlying $\dot{V}O_{2max}$ and/or relative oxygen transport capability. In addition, the included non-gas-analyzed measures of CRF authentically imitate activities of daily living (e.g., running, cycling, stair climbing) and are suggestive of declines in functional and exercise capability.

Physiological changes are in turn affected by physical changes such as increased fatness, as well as behavioral changes such as decreased physical activity levels or increased sedentary time, which may result in reduced cardiovascular function [3, 27, 39]. Consider first increases in fatness, for which an international increase is well established [35, 44]. There are both mechanistic and correlational arguments to support why increases in fatness will influence declines in CRF. First, because fat constitutes an additional load, increased fatness will decrease relative $\dot{V}O_{2max}$ (ml/kg/min) approximately on a pro rata basis [45, 46], and in contrast to a passive load, it will result in additional metabolic maintenance costs (e.g., breathing, thermoregulation) [47]. However, while increased fatness will increase the energy demand, it is probable that large increases in fatness are required to meaningfully change energy requirements [48]. Second, increases in fatness have coincided with declines in CRF. Consistent with evidence in children [32, 49], this study found that increased country-level adult obesity was very strongly and negatively correlated with country-level decreased adult CRF. Assuming the observed ecological correlation is causal, this temporal connection suggests that obesity prevention strategies (e.g., reduced energy intake relative to expenditure, increased energy expenditure) might be a suitable population approach to slowing the decline in CRF, or that strategies to improve or maintain CRF (e.g., increased moderate-to-vigorous physical activity) might help to slow the increase in adult obesity. Improving or maintaining CRF may be relatively more important, or at least as important, for cardiovascular disease risk and prognosis than the prevention of obesity [51, 52]. Third, using a matching analysis (where people tested at one time point are matched for sex, age, and fatness with people tested years later), increases in fatness have been estimated to explain 35–70% of the declines in




Fig. 3 National temporal trends in mean cardiorespiratory fitness (CRF) at the age and sex level between 1967 and 2016. Data are standardized to the year 1990 = 100%, with higher values (>100%) indicating better CRF and negative values (<100%) indicating better CRF and negative values (<100%) indicating poorer CRF. The black lines represent the national changes in mean CRF for females (♀; left panels) and those aged < 40 years (right panels), the grey lines represent the national changes in mean CRF for males (♂; left panels) and those aged ≥40 years (right panels), with upward sloping lines indicating increases over time and downward sloping lines indicating declines over time. Mean (95% CI) percent changes (per decade) are shown at the top of each panel

CRF [50, 53]. Unfortunately, the residual variability remains unexplained. Importantly, recent increases in adult fatness have coincided with increases in fat-free mass [54], both of which affect the balance between energy supply and energy demand. While increased fat-free mass will improve the metabolic potential of the exercising muscles (e.g., improved capacity for glycogen storage, increased muscular strength resulting from increased cross-sectional area, increased number of mitochondria), such a benefit to energy supply will likely be outweighed by the increase in energy demand resulting from increased fatness.

Of course temporal trends in behaviors are also probably involved [27, 28], which are in turn likely influenced by broad socioeconomic and/or environmental trends [27, 28]. Because of the difficulty in obtaining accurate measurements and temporal differences in sampling and methodology, trend data on adult physical activity levels are scarce. At this time there is no compelling evidence for international increases in physical activity levels or declines in sedentary behaviors [55, 56]. Adult trend data from high-income countries suggest that leisure-time physical activity has increased [57–59] and occupational physical activity has declined [58, 60, 61]. Similarly in children and adolescents, physical activity levels and sport participation appear to have remained stable [62, 63] or increased [64]. Because CRF assesses an individual’s physiological response to their total physical activity profile, declines in adult CRF are therefore suggestive of declines in overall physical activity levels. Furthermore, while declines in CRF were meaningful correlates of trends in socioeconomic (HDI) and environmental (air pollution and urbanization) indicators (Table 3), our confidence in these correlations was low due to: (a) the small number of countries resulting in reduced statistical power, (b) the short time window over which trend data were available (e.g., post-1990 for HDI and air pollution), and (c) the homogeneity in the available trend data between countries.

Table 3 Potential correlates of the trends in adult cardiorespiratory fitness (CRF)

| Variable | Data source | Description | Correlation (95% CI) |
|------------------------------------|--|---|------------------------|
| Prevalence of obesity (%) | Ng et al. [35] Trend data available for 7 of 8 (88%) countries between 1990 and 2013 | Calculated as the change (per decade) in mean country-level prevalence of men and women aged ≥ 20 years (age standardized) classified as obese based on the International Obesity Task Force definition. A positive change indicated an increase in the mean prevalence of obese and a negative change indicated a decline | -0.77 (-0.96 to -0.03) |
| Human development index (HDI) | The United Nations [36] Trend data available for 7 of 8 (88%) countries between 1990 and 2015 | Calculated as the change (per decade) in mean country-level achievement in a variety of indicators related to standards of living. A positive change indicated an increase in the mean standard of living and a negative change indicated a decline | 0.15 (-0.68 to 0.81) |
| Urbanization (%) | World Bank [37] Trend data available for 8 of 8 (100%) countries between 1967 and 2016 | Calculated as the change (per decade) in the percentage of people living in urban areas. A positive change indicated an increase in urbanization and a negative change indicated a decline | -0.50 (-0.89 to 0.32) |
| Air pollution (PM _{2.5}) | World Bank [37] Trend data available for 7 of 8 (88%) countries between 1990 and 2015 | Calculated as the change (per decade) in population-weighted annual mean exposure to ambient PM _{2.5} pollution (i.e., exposure of a population to concentrations of suspended particles measuring <2.5 microns in aerodynamic diameter). A positive change indicated an increase in mean air pollution exposure and a negative change indicated a decline | 0.15 (-0.68 to 0.81) |

In order to improve CRF, the World Health Organization [65] recommends that adults participate in at least 150 min of moderate intensity aerobic exercise (continuous, rhythmic, dynamic, large muscle group exercise such as walking, running, cycling, swimming) each week, or at least 75 min of vigorous intensity aerobic exercise each week, or an equivalent combination thereof. For additional health benefits, adults are encouraged to double the recommended weekly dose [65]. The World Health Organization’s recent global action plan on physical activity [66], which targets a 15% relative reduction in the global prevalence of physical inactivity in adults and adolescents by 2030, provides a comprehensive roadmap for global efforts to increase the physical activity levels of all people. The monitoring of CRF—a proximal outcome of total physical activity levels—might be a favorable approach to complement the existing global action plan [67]. Globally, physical inactivity is estimated to cost INT\$54 billion per year in direct health care, with an additional INT\$14 billion attributable to lost productivity [68].

4.2 Temporal Differences in CRF Between the Sexes and Different Age Groups

Figure 3 (two bottom panels) indicates that the collective decline in mean CRF was larger for men than for women and for young adults than for middle-aged adults. These temporal differences have previously been identified in adults [40, 69] and children [29]. Ekblom et al. [40] found that over the period 1990–2000 median CRF ($\dot{V}O_{2\max}$ in ml/kg/min) declined in men but not women, and declined with increasing age in both men and women. Similarly, Willis et al. [69] found that while mean CRF ($\dot{V}O_{2\max}$ in ml/kg/min estimated from maximal treadmill walking/running) increased in 52,785 men aged 20–74 years between 1970 and 2009, CRF declined somewhat after 2000, with the largest declines observed in young men aged 20–34 years. In our recent analysis of 965,264 children aged 9–17 years from 19 high-income and upper-middle-income countries between 1981 and 2014, we found that the decline in mean CRF (ml/kg/min estimated from 20-m shuttle run test performance) was larger for boys than for girls [29]. Coupled with evidence indicating that improving CRF at a younger age confers the greatest mortality benefit (e.g., each 1 MET increase in CRF is associated with a 15% decrease in mortality risk in men <60 years as opposed to an 11% decrease in mortality risk in men ≥ 60 years [70]), these trend data highlight the importance of targeting exercise and/or physical activity interventions to improve or maintain CRF across the lifespan, especially in early adulthood.

These temporal differences are difficult to explain in the absence of concurrent data collected on the same adults as those examined in this study. Data from the International Obesity Task Force [35] indicate that recent increases in age-standardized adult obesity (in the countries included in this study at least) have been somewhat larger for men (~ 1.7% per decade) than for women (~ 1.1% per decade), and for young adults (~ 1.6% per decade) than for middle-aged adults (~ 1.2% per decade). Similarly, Ekblom et al. [40] found that both BMI and the prevalence of overweight and obesity increased in men aged 20–59 years, with the largest increases found in young men aged 20–29 years; no changes were found in women. While the temporal coincidence of these patterns is potentially circumstantial, it does at least suggest a strong association between fatness and CRF, although the direction of the causal relationship is unclear.

It is also possible that these sex-related temporal differences in CRF are due to temporal differences in physical activity participation [29]. While good physical activity trend data are hard to identify, data on participation in elite sport and occupational physical activity and use of time indicate that gender differences exist. For example, female participation at the Olympic Games has steadily risen since its inception in 1900, with females comprising 45% of all athletes in 2016, increasing from 26% in 1988 and 11% in 1960 [71]. This diminishing gender gap in elite sport may reflect that international sport participation has increased for all females. There is also evidence of a gender difference in trends in occupational physical activity and use of time in some [72, 73] but not all [74, 75] countries. For example, in men from Sweden and the USA, occupational physical activity and time spent in paid work has declined, whereas in women, there has been no corresponding change in occupational physical activity and a slight increase in time spent in paid work [72, 73]. Temporal trends in occupational physical activity (e.g., increased automation and computerization of labor) may have therefore affected the CRF of men over time more so than that of women. Perhaps the recent promotion of gender equality policies and programs that aim to empower women and girls [76] has helped to reduce the gender gap in CRF through leisure-time and occupational physical activity. Sex-related differences in motivation levels may also be involved.

| Year of testing | Canada | China | Finland | Japan | Norway | Poland | Republic of Korea | Singapore |
|-----------------|--------|-------|---------|-------|--------|--------|-------------------|-----------|
| 1967 | | | | • | | | | |
| 1968 | | | | • | | | | |
| 1969 | | | | • | | | | |
| 1970 | | | | • | | | | |
| 1971 | | | | • | | | | |
| 1972 | | | | • | | | | |
| 1973 | | | | • | | | | |
| 1974 | | | | • | | | | |
| 1975 | | | • | • | | | | |
| 1976 | | | • | • | | | | |
| 1977 | | | • | • | | | | |
| 1978 | | | • | • | | | | |
| 1979 | | | • | • | | • | | |
| 1980 | | | • | • | • | | | |
| 1981 | • | | • | • | • | | | • |
| 1982 | | | • | • | • | | | |
| 1983 | | | • | • | • | | | |
| 1984 | | | • | • | • | | | |
| 1985 | | • | • | • | • | | | |
| 1986 | | | • | • | | | | |
| 1987 | | | • | • | | | | |
| 1988 | | | • | • | | | • | |
| 1989 | | | • | • | | • | • | |
| 1990 | | | • | • | | | • | |
| 1991 | | • | • | • | | | | • |
| 1992 | | | • | • | | | • | |
| 1993 | | | • | • | | | | |
| 1994 | | | • | • | | | | |
| 1995 | | • | • | • | | | • | |
| 1996 | | | • | • | | | | |
| 1997 | | | • | • | | | | |
| 1998 | | | • | • | | | • | |
| 1999 | | | • | • | | • | | |
| 2000 | | • | • | • | | | | |
| 2001 | | | • | • | | | • | |
| 2002 | | | • | • | • | | | |
| 2003 | | | • | • | | | | |
| 2004 | | | • | • | | | • | |
| 2005 | | • | • | • | | | | |
| 2006 | | | • | • | | | | |
| 2007 | | | • | • | | | • | |
| 2008 | | | • | • | | | | |
| 2009 | • | | • | • | | | | |
| 2010 | | • | • | • | | | | |
| 2011 | | | • | • | | | | |
| 2012 | | | • | • | | | | |
| 2013 | | | • | • | | | | |
| 2014 | | • | • | • | | | | |
| 2015 | | | • | • | | | | |
| 2016 | | | • | • | | | | |

Table 2 Distribution of national surveys from which temporal trends in cardiorespiratory fitness were calculated

4.3 Strengths and Limitations

This study systematically reviewed and analyzed data from eight unique datasets/studies to estimate international and national temporal trends in the CRF of 2,525,827 adults between 1967 and 2016. The inclusion of only population representative national temporal data on adults who were tested using validated measures of CRF, coupled with the use of weighted regression and a post-stratification population weighting procedure, resulted in high confidence that the reported declines are in fact real. A sensitivity analysis showed that the removal of countries (China, Japan, and the Republic of Korea) with very large population sizes ($n > 50,000,000$) and the highest numbers of country-sex-age-year groups had a negligible effect ($ES < 0.1$) on the collective trend in CRF, providing support that the reported international trend in CRF was not biased by these countries.

While this is the first systematic analysis of temporal trends in adult CRF, it used a detailed statistical approach that has been previously adopted in similar reviews on children's CRF [27–29], thus widening the lens on the international trends picture and allowing for direct comparisons between children and adults. For example, Fig. 4 shows the international temporal trends in mean CRF for children and adults from high- and upper-middle-income countries between 1981 and 2014. Comparative trend data were taken from Tomkinson et al.

[29]. Examination of Fig. 4 reveals that there has been a moderate decline in CRF for both children ($ES - 0.71$) and adults ($ES - 0.64$) since 1981, although the shapes of the temporal trends have differed. In children, the rate of decline slowed from 1981 to 2000 and stabilized near zero thereafter; in adults, the rate of decline progressively increased with every decade. These data suggest that the current decline in adult CRF might diminish in subsequent decades when today's children (whose CRF levels have remained reasonably stable for the past decade and a half) become adults. Alternatively, because the transition from adolescence to adulthood represents a period of significant change where activities of daily living are restructured, these temporal differences might indicate a widening of the gap in CRF levels between adolescence and adulthood over time, resulting in a relatively larger decline in CRF in early adulthood. While the underlying reasons for these age-related temporal differences are not clear, as with temporal differences between men and women and young and middle-aged adults, differential trends in fatness, physical activity, and sedentary behaviors are probably involved. Data from Olds et al. [77] indicate that childhood overweight and obesity has stabilized since the mid-to-late 1990s in some high-income countries, which may help explain the recent plateau in children's CRF.




Fig. 4 International temporal trends in mean cardiorespiratory fitness (CRF) for children (9–17 years) and adults (18–59 years) from high and upper-middle-income countries between 1981 and 2014. Data are standardized to the year 1990=100%, with higher values (>100%) indicating better CRF and negative values (<100%) indicating poorer CRF. The black trend line represents the international change in mean CRF for adults (18–59 years) and the grey trend line the international change in mean CRF for children (9–17 years), with upward sloping lines indicating increases over time and downward sloping lines indicating declines over time; trend data for children are from Tomkinson et al. [29]

It is possible that the country-level CRF data have not always been collected under precisely the same conditions, resulting in systematic differences in testing conditions (e.g., climate, altitude, practice, and running surfaces) and measurement errors (e.g., methodological drift and diurnal variation). However, the inclusion of only large randomized national fitness survey data would have minimized sampling- and methodology-related issues, meaning the reported trends were unlikely to be systematically biased. This study was also limited to the examination of temporal trends in mean values, which could be systematically biased if concomitant trends in skewness occurred. Consistent with data on children [49, 78], Santtila et al. [79] indicated that the decline in the 12-min distance run performance of 20-year-old Finnish men was skewed towards the low fitness end of the distribution. On the one hand, this suggests that the reported declines in mean CRF are inflated; on the other hand, the trends in skewness reported by Santtila et al. [79] are the likely artifact of expressing CRF as the distance run in 12 min.

It is also important to remember that the reported trends are representative of only seven very high human development (high-income) countries and one high human development (upper-middle-income) country, and it is not known whether similar trends exist for other high- and upper-middle-income countries or for low-income countries that may be experiencing a physical activity transition [80]. In addition, the reported trends were estimated from available country-sex-age-specific trend data, which may not be representative of trends across all adults within the included countries or all measurement years. The broad data coverage of the large Japanese dataset (representing 55% of all data points) meant that the international trend in CRF was largely driven by data from a single country, specifically in the first decade over which only Japanese data were available. Finally, given that the international trends in CRF were estimated using different CRF tests, each of which imposes different physiological and psychosocial demands, it could be that the trends (percent and standardized) for one test do not equate to the trends for another. It is likely, however, that the reported trends are largely reflective of declines in construct CRF because the included tests demonstrate moderate-to-very high criterion validity and provide a strong aerobic physiological challenge. Nonetheless, the reported declines reflect declines in exercise/functional capability.

5 Conclusion

This study indicates that there has been a meaningful international decline in the CRF of adults since 1980, which has progressively increased in magnitude every decade thereafter. Declines in adult CRF were experienced by all included countries, with declines larger for men than for women, and for young adults than for middle-aged adults. Although it is not known why the sex- and age-related temporal differences exist, temporal differences in adult obesity and physical activity levels may be involved. Another key finding was that country-level declines in adult CRF coincided with increases in adult obesity, with the smallest national declines in CRF observed in countries with the smallest increases in obesity. Given the etiology of the decline in CRF is of more than just theoretical interest, it is important that trends in CRF and other cardiovascular disease risk factors (e.g., obesity, physical inactivity, sedentary behavior) be tracked if we want to develop appropriate strategies to reverse the current decline in CRF. Consistent with CRF data in children and adolescents, there are currently no available trend data for adults from low- and middle-income countries. In an effort to enhance knowledge, we advocate for the development of continuous national and international surveillance health systems (e.g., through web-surveillance of directly-measured and estimated CRF [81, 82]), especially among low- and middle-income countries, in order to monitor health and fitness and to guide national and international action. The inclusion of measures of CRF in standardized health surveillance systems would: (a) provide a robust, standardized alternative to assess country-level population health, (b) complement existing health measures (e.g., physical activity, obesity) to provide more meaningful insight into population health, (c) better inform practitioners and policy makers about the impact of health promotion policies and interventions, and (d) help predict future health and guide public health resource allocation [81].

Data availability statement The datasets analyzed in this review are available from the corresponding author on reasonable request.

Acknowledgements We would like to thank the authors of the included studies for generously clarifying details of their studies and/or for providing additional data. We would also like to thank Dr. Dae-Yeon Kim, Dr. Yang Liu, Dr. Tetsuhiro Kidokoro, Dr. Shingo Noi, and Dr. Koya Suzuki for their assistance with the national fitness data from China, Japan and the Republic of Korea.

Author contributions GRT and NRL developed the research question and designed the study. GRT and NRL had full access to all the data in the study and take responsibility for the integrity of the data. GRT and NRL led the statistical analysis, synthesis of results, and writing of the report. All authors contributed to the interpretation of results, editing and critical reviewing of the final report, and approved the final report.

Compliance with Ethical Standards

Conflict of interest Nicholas R. Lamoureux, John S. Fitzgerald, Kevin I. Norton, Todd Sabato, Mark S. Tremblay, and Grant R. Tomkinson declare that they have no conflict of interest. Funding No funding was received for this project.

References

1. Ross R, Blair SN, Arena R, Church TS, Després J, Franklin BA, et al. Importance of assessing cardiorespiratory fitness in clinical practice: a case for fitness as a clinical vital sign: a scientific statement from the American Heart Association. *Circulation*. 2016;134(24):e653–99.

2. Harber MP, Kaminsky LA, Arena R, Blair SN, Franklin BA, Myers J, et al. Impact of cardiorespiratory fitness on all-cause and disease-specific mortality: advances since 2009. *Prog Cardiovasc Dis*. 2017;60(1):11–20.
3. Armstrong N, Tomkinson G, Ekelund U. Aerobic fitness and its relationship to sport, exercise training and habitual physical activity during youth. *Br J Sports Med*. 2011;45(11):849–58.
4. Fletcher GF, Blair SN, Blumenthal J, Caspersen C, Chaitman B, Epstein S, et al. Statement on exercise. Benefits and recommendations for physical activity programs for all Americans. A statement for health professionals by the Committee on Exercise and Cardiac Rehabilitation of the Council on Clinical Cardiology, American Heart Association. *Circulation*. 1992;86(1):340–4.
5. Blair SN, Kohl HW 3rd, Paffenbarger RS Jr, Clark DG, Cooper KH, Gibbons LW. Physical fitness and all-cause mortality: a prospective study of healthy men and women. *JAMA*. 1989;262(17):2395–401.
6. Berry JD, Willis B, Gupta S, Barlow CE, Lakoski SG, Khera A, et al. Lifetime risks for cardiovascular disease mortality by cardiorespiratory fitness levels measured at ages 45, 55, and 65 years in men The Cooper Center Longitudinal Study. *J Am Coll Cardiol*. 2011;57(15):1604–10.
7. Kodama S, Saito K, Tanaka S, Maki M, Yachi Y, Asumi M, et al. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *JAMA*. 2009;301(19):2024–35.
8. Lakka TA, Venäläinen JM, Rauramaa R, Salonen R, Tuomilehto J, Salonen JT. Relation of leisure-time physical activity and cardiorespiratory fitness to the risk of acute myocardial infarction in men. *N Engl J Med*. 1994;330(22):1549–54.
9. Lee CD, Blair SN. Cardiorespiratory fitness and stroke mortality in men. *Med Sci Sports Exerc*. 2002;34(4):592–5.
10. Lee DC, Sui X, Ortega FB, Kim YS, Church TS, Winett RA, et al. Comparisons of leisure-time physical activity and cardiorespiratory fitness as predictors of all-cause mortality in men and women. *Br J Sports Med*. 2011;45(6):504–10.
11. McAuley P, Pittsley J, Myers J, Abella J, Froelicher VF. Fitness and fatness as mortality predictors in healthy older men: the Veterans Exercise Testing Study. *J Gerontol A Biol Sci Med Sci*. 2009;64(6):695–9.
12. Sui X, LaMonte MJ, Laditka JN, Hardin JW, Chase N, Hooker SP, et al. Cardiorespiratory fitness and adiposity as mortality predictors in older adults. *JAMA*. 2007;298(21):2507–16.
13. Crump C, Sundquist J, Winkleby MA, Sieh W, Sundquist K. Physical fitness among Swedish military conscripts and long-term risk for type 2 diabetes mellitus: a cohort study. *Ann Intern Med*. 2016;164(9):577–84.
14. Jensen MT, Holtermann A, Bay H, Gyntelberg F. Cardiorespiratory fitness and death from cancer: a 42-year follow-up from the Copenhagen Male Study. *Br J Sports Med*. 2017;51(18):1364–9.
15. Park MS, Chung SY, Chang Y, Kim K. Physical activity and physical fitness as predictors of all-cause mortality in Korean men. *J Korean Med Sci*. 2009;24(1):13–9.

16. Robsahm TE, Falk RS, Heir T, Sandvik L, Vos L, Erikssen JE, et al. Measured cardiorespiratory fitness and self-reported physical activity: associations with cancer risk and death in a long-term prospective cohort study. *Cancer Med*. 2016;5:2136–44.
17. Sawada SS, Lee IM, Naito H, Kakigi R, Goto S, Kanazawa M, et al. Cardiorespiratory fitness, body mass index, and cancer mortality: a cohort study of Japanese men. *BMC Public Health*. 2014;14:1012.
18. Carnethon MR, Evans NS, Church TS, Lewis CE, Schreiner PJ, Jacobs DR Jr, et al. Joint associations of physical activity and aerobic fitness on the development of incident hypertension: coronary artery risk development in young adults. *Hypertension*. 2010;56(1):49–55.
19. Carnethon MR, Gidding SS, Nehgme R, Sidney S, Jacobs DR Jr, Liu K. Cardiorespiratory fitness in young adulthood and the development of cardiovascular disease risk factors. *JAMA*. 2003;290(23):3092–100.
20. Gill JM, Malkova D. Physical activity, fitness and cardiovascular disease risk in adults: interactions with insulin resistance and obesity. *Clin Sci (Lond)*. 2006;110(4):409–25.
21. Gupta S, Rohatgi A, Ayers CR, Willis BL, Haskell WL, Khera A, et al. Cardiorespiratory fitness and classification of risk of cardiovascular disease mortality. *Circulation*. 2011;123(13):1377–83.
22. Holtermann A, Marott JL, Gyntelberg F, Søgaard K, Mortensen OS, Prescott E, et al. Self-reported cardiorespiratory fitness: prediction and classification of risk of cardiovascular disease mortality and longevity—a prospective investigation in the Copenhagen City Heart Study. *J Am Heart Assoc*. 2015;4(1):e001495.
23. Laukkanen JA, Rauramaa R, Salonen JT, Kurl S. The predictive value of cardiorespiratory fitness combined with coronary risk evaluation and the risk of cardiovascular and all-cause death. *J Intern Med*. 2007;262(2):263–72.
24. Myers J, Nead KT, Chang P, Abella J, Kokkinos P, Leeper NJ. Improved reclassification of mortality risk by assessment of physical activity in patients referred for exercise testing. *Am J Med*. 2015;128(4):396–402.
25. Blair SN. Physical inactivity: the biggest public health problem of the 21st century. *Br J Sports Med*. 2009;43(1):1–2.
26. Tomkinson GR, Olds TS. Field tests of fitness. In: Armstrong N, Van Mechelen W, editors. *Paediatric exercise science and medicine*. 2nd ed. Oxford: Oxford University Press; 2008. p. 109–28.
27. Tomkinson GR, Léger LA, Olds TS, Cazorla G. Secular trends in the performance of children and adolescents (1980–2000): an analysis of 55 studies of the 20 m shuttle run test in 11 countries. *Sports Med*. 2003;33(4):285–300.
28. Tomkinson GR, Olds TS. Secular changes in pediatric aerobic fitness test performance: the global picture. *Med Sport Sci*. 2007;50:46–66.
29. Tomkinson GR, Lang JJ, Tremblay MS. Temporal trends in the cardiorespiratory fitness of children and adolescents representing 19 high-income and upper middle-income countries between 1981 and 2014. *Br J Sports Med*. 2017. <https://doi.org/10.1136/bjsports-2017-097982> (Epub ahead of print).
30. Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev*. 2015;4:1.

31. Leger L. Aerobic performance. In: Docherty D, editor. *Measurement in pediatric exercise science*. Champaign: Human Kinetics; 1996. p. 183–223.
32. Tomkinson GR, Macfarlane D, Noi S, Dae-Yeon K, Wang Z, Hong R. Temporal changes in long-distance running performance of Asian children between 1964 and 2009. *Sports Med*. 2012;42(4):267–79.
33. Levy PS, Lemeshow S. Stratification random sampling: further issues. In: Levy PS, Lemeshow S, editors. *Sampling of populations: methods and application*. Hoboken: Wiley; 2008. p. 143–88.
34. United Nations, Department of Economic and Social Affairs, Population Division. *World population prospects: the 2017 revision, key findings and advance tables*. Working Paper. ESA/P/WP/248. New York: United Nations; 2017.
35. Ng M, Fleming T, Robinson M, Thomson B, Graetz N, Margono C, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 2014;384(9945):766–81.
36. United Nations Development Programme. *Human Development Report 2016: human development for everyone*. New York: United Nations Development Programme; 2016.
37. World Bank. *DataBank: World development indicators*. <http://databank.worldbank.org/data/reports.aspx?source=world-development-indicators>. Accessed 13 Aug 2018.
38. Craig CL, Shields M, Leblanc AG, Tremblay MS. Trends in aerobic fitness among Canadians, 1981 to 2007–2009. *Appl Physiol Nutr Metab*. 2012;37(3):511–9.
39. Dyrstad SM, Berg T, Tjelta LI. Secular trends in aerobic fitness performance in a cohort of Norwegian adolescents. *Scand J Med Sci Sports*. 2012;22(6):822–7.
40. Ekblom B, Engström L, Ekblom O. Secular trends of physical fitness in Swedish adults. *Scand J Med Sci Sports*. 2007;17(3):267–73.
41. Cink RE, Thomas TR. Validity of the Åstrand-Ryhming nomogram for predicting maximal oxygen intake. *Br J Sports Med*. 1981;15(3):182–5.
42. Mayorga-Vega D, Bocanegra-Parrilla R, Ornelas M, Viciania J. Criterion-related validity of the distance- and time-based walk/ run field tests for estimating cardiorespiratory fitness: a systematic review and meta-analysis. *PLoS One*. 2016;11(3):e0151671.
43. Weller IM, Thomas SG, Gledhill N, Paterson D, Quinney A. A study to validate the modified Canadian Aerobic Fitness Test. *Can J Appl Physiol*. 1995;20(2):211–21.
44. NCD Risk Factor Collaboration (NCD-RisC). Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19.2 million participants. *Lancet*. 2016;387(10026):1377–96.
45. Cureton KJ, Sparling PB, Evans BW, Johnson SM, Kong UD, Purvis JW. Effect of experimental alterations in excess weight on aerobic capacity and distance running performance. *Med Sci Sports*. 1978;10(3):194–9.
46. Goran M, Fields DA, Hunter GR, Herd SL, Weinsier RL. Total body fat does not influence maximal aerobic capacity. *Int J Obes Relat Metab Disord*. 2000;24(7):841–8.

47. Olds TS, Ridley K, Tomkinson GR. Declines in aerobic fitness: are they only due to increasing fatness? *Med Sport Sci.* 2007;50:226–40.
48. Cureton KJ, Boileau RA, Lohman TG, Misner JE. Determinants of distance running performances in children: analysis of a path model. *Res Q.* 1977;48(2):270–9.
49. Dollman J, Olds T, Norton K, Stuart D. The evolution of fitness and fatness in 10–11-year-old Australian schoolchildren: changes in distributional characteristics between 1985 and 1997. *Pediatr Exerc Sci.* 1999;11(2):108–21.
50. Olds T, Dollman J. Are changes in distance-run performance of Australian children between 1985 and 1997 explained by changes in fatness? *Pediatr Exerc Sci.* 2004;16(3):201–9.
51. Kennedy AB, Lavie CJ, Blair SN. Fitness or fatness: which is more important? *JAMA.* 2018;319(3):231–2.
52. Oktay AA, Lavie CJ, Kokkinos PF, Parto P, Pandey A, Ventura HO. The interaction of cardiorespiratory fitness with obesity and the obesity paradox in cardiovascular disease. *Prog Cardiovasc Dis.* 2017;60(1):30–44.
53. Albon HM, Hamlin MJ, Ross JJ. Secular trends and distributional changes in health and fitness performance variables of 10–14-year-old children in New Zealand between 1991 and 2003. *Br J Sports Med.* 2010;44(4):263–9.
54. Knapik JJ, Sharp MA, Steelman RA. Secular trends in the physical fitness of United States Army recruits on entry to service, 1975–2013. *J Strength Cond Res.* 2017;31(7):2030–52.
55. Hallal PC, Andersen LB, Bull FC, Guthold R, Haskell W, Ekelund U, et al. Global physical activity levels: surveillance progress, pitfalls, and prospects. *Lancet.* 2012;380(9838):247–57.
56. Sallis JF, Bull F, Guthold R, Heath GW, Inoue S, Kelly P, et al. Progress in physical activity over the Olympic quadrennium. *Lancet.* 2016;388(10051):1325–36.
57. Juneau CE, Potvin L. Trends in leisure-, transport-, and work-related physical activity in Canada 1994–2005. *Prev Med.* 2010;51(5):384–6.
58. Knuth AG, Hallal PC. Temporal trends in physical activity: a systematic review. *J Phys Act Health.* 2009;6(5):548–59.
59. Devonshire-Gill KR, Norton KI. Australian adult physical activity sufficiency trend data: positive, prevalent, and persistent changes 2002–2012. *J Phys Act Health.* 2018;15(2):117–26.
60. Brownson RC, Boehmer TK, Luke DA. Declining rates of physical activity in the United States: what are the contributors? *Annu Rev Public Health.* 2005;26:421–43.
61. Stamatakis E, Chaudhury M. Temporal trends in adults' sports participation patterns in England between 1997 and 2006: the Health Survey for England. *Br J Sports Med.* 2008;42(11):901–8.
62. Booth VM, Rowlands AV, Dollman J. Physical activity temporal trends among children and adolescents. *J Sci Med Sport.* 2015;18(4):418–25.
63. Ekelund U, Tomkinson G, Armstrong N. What proportion of youth are physically active? Measurement issues, levels and recent time trends. *Br J Sports Med.* 2011;45(11):859–65.

64. Kalman M, Inchley J, Sigmundova D, Iannotti RJ, Tynjälä JA, Hamrik Z, et al. Secular trends in moderate-to-vigorous physical activity in 32 countries from 2002 to 2010: a cross-national perspective. *Eur J Public Health*. 2015;25(S2):37–40.
65. World Health Organization. Global recommendations on physical activity for health. Geneva: World Health Organization; 2010.
66. World Health Organization. Global action plan on physical activity 2018–2030: more active people for a healthier world. Geneva: World Health Organization; 2018.
67. Lang JJ, Wolfe Phillips E, Orpana HM, Tremblay MS, Ross R, Ortega FB, et al. Field-based measurement of cardiorespiratory fitness as a way to evaluate physical activity interventions in very high-, high-, middle-, and low-income countries. *Bull World Health Organ*. 2018;96:794–6.
68. Ding D, Lawson KD, Kolbe-Alexandar TL, Finkelstein EA, Katzmarzyk PT, Mechelen W, et al. The economic burden of physical inactivity: a global analysis of major non-communicable diseases. *Lancet*. 2016;388(10051):1311–24.
69. Willis BL, Morrow JR Jr, Jackson AW, Defna LF, Cooper KH. Secular change in cardiorespiratory fitness of men: Cooper Center Longitudinal Study. *Med Sci Sports Exerc*. 2011;43(11):2134–9.
70. Kokkinos P, Faselis C, Myers J, Sui X, Zhang J, Blair SN. Age-specific exercise capacity threshold for mortality risk assessment in male veterans. *Circulation*. 2014;130(8):653–8.
71. International Olympic Committee. IOC factsheet: women in the Olympic movement—update January 2016. https://stillmed.olympic.org/Documents/Reference_documents/Factsheets/Women_in_Olympic_Movement.pdf. Accessed 28 Mar 2018.
72. Lindahl B, Stegmayr B, Johansson I, Weinehall L, Hallmans G. Trends in lifestyle 1986–99 in a 25- to 64-year-old population of the Northern Sweden MONICA project. *Scand J Public Health*. 2003;31(S61):31–7.
73. Robinson JP, Godbey G. Time for life: the surprising ways Americans use their time. University Park: Pennsylvania State University Press; 1999.
74. Simmons G, Jackson R, Swinburn B, Yee RL. The increasing prevalence of obesity in New Zealand: is it related to recent trends in smoking and physical activity? *N Z Med J*. 1996;109(1018):90–2.
75. Stamatakis E, Ekelund U, Wareham NJ. Temporal trends in physical activity in England: the Health Survey for England 1991 to 2004. *Prev Med*. 2007;45(6):416–23.
76. United Nations. Sustainable Development Goals: 17 goals to transform our world. Goal 5: Achieve gender equality and empower all women and girls. <http://www.un.org/sustainabledevelopment/gender-equality/>. Accessed 28 Mar 2018.
77. Olds T, Maher C, Zumin S, Péneau S, Lioret S, Castetbon K, et al. Evidence that the prevalence of childhood overweight is plateauing: data from nine countries. *Int J Pediatr Obes*. 2011;6(5–6):342–60.
78. McNaughton L, Morgan R, Smith P, Hannan G. An investigation into the fitness levels of Tasmanian primary schoolchildren. *ACHPER Healthy Lifestyles J*. 1996;43(1):4–10.
79. Santtila M, Kyröläinen H, Vasankari T, Tiainen S, Palvalin K, Häkkinen A, et al. Physical fitness profiles in young Finnish men during the years 1975–2004. *Med Sci Sports Exerc*. 2006;38(11):1990–4.

80. Katzmarzyk PT, Mason C. The physical activity transition. *J Phys Act Health*. 2009;6(3):269–80.
81. Lang JJ, Tomkinson GR, Janssen I, Ruiz JR, Ortega FB, Léger L, et al. Making a case for cardiorespiratory fitness surveillance among children and youth. *Exerc Sport Sci Rev*. 2018;46(2):66–75.
82. Nauman J, Tauschek LC, Kaminsky LA, Nes BM, Wisløf U. Global fitness levels: findings from a web-based surveillance report. *Prog Cardiovasc Dis*. 2017;60(1):78–88.