



2017

## Opioid Maintenance Treatment: Methadone, Buprenorphine, and Naltrexone

Nicole Lemieux  
*University of North Dakota*

Follow this and additional works at: <https://commons.und.edu/pas-grad-posters>

 Part of the [Substance Abuse and Addiction Commons](#)

[How does access to this work benefit you? Let us know!](#)

---

### Recommended Citation

Lemieux, Nicole, "Opioid Maintenance Treatment: Methadone, Buprenorphine, and Naltrexone" (2017).  
*Physician Assistant Scholarly Project Posters*. 44.  
<https://commons.und.edu/pas-grad-posters/44>

This Poster is brought to you for free and open access by the Department of Physician Studies at UND Scholarly Commons. It has been accepted for inclusion in Physician Assistant Scholarly Project Posters by an authorized administrator of UND Scholarly Commons. For more information, please contact [und.common@library.und.edu](mailto:und.common@library.und.edu).



# Opioid Maintenance Treatment: Methadone, Buprenorphine, and Naltrexone

*Nicole Lemieux, PA-S*  
*Department of Physician Assistant Studies, University of North Dakota School of Medicine & Health Sciences*  
*Grand Forks, ND 58202-9037*



## Abstract

Opioid dependence plagues the United States with millions of people suffering from the effects. Throughout recent years, there has been a surge in research in treatment options for patients suffering from opioid dependence. Three medications have stood out: methadone, buprenorphine, and naltrexone. The purpose of this study was to investigate the three and decipher which provides the best outcomes for patients. Research was conducted by reviewing literature via PubMed, PsychInfo and Cochrane. A review of the literature found that methadone has been used the longest and has been shown to be efficacious, but can be severely limiting for patients due to lack of clinics and the need for daily dosings. As a result, buprenorphine was developed as an alternative medication with similar efficacy and retention rates to combat the daily dosages and provide a more readily available medication. Unfortunately buprenorphine also has geographical limitations. Naltrexone is the newest medication added to the mix. Orally, it has poor retention, but an extended release form has been developed that is showing promise. There are no geographical limitations associated with naltrexone and the extended release form allows for only monthly injections. Going forward with opioid treatment, medications like naltrexone and those that are more readily available will become more widely used for treatment as patient preference is pointing towards medications that are not limited by geography or frequent dosing.

## Introduction

- Opioid addiction plagues our country. Short term opioid use has been shown to be beneficial for patients, but many of those patients continue to receive opioids for years (Lembke, Humphreys, & Newmark, 2016). The addiction ranges from patients receiving chronic opioid prescriptions to those buying them on the street and abusing. This epidemic has become a significant public health problem. There is increased morbidity and mortality associated with opioid use, as well as an increase in the spread of HIV and hepatitis (Nielsen et al., 2016).
- Opioid maintenance therapy has been shown to reduce the use of abused opioids. Patients are able to return to normal social functioning and decrease their criminal activity. This also decreases the transmission of infectious diseases such as HIV and hepatitis (Nunes et al.,2015).
- The Substance Abuse and Mental Health Services Administration endorses three medications to use for opioid maintenance treatment: buprenorphine, methadone, and naltrexone. While all three have their benefits and their challenges, this researcher has reviewed the current literature to make an informed decision as to which is the best treatment option of the three (buprenorphine, methadone, and naltrexone) for patients in need of opioid maintenance treatment.

## Statement of the Problem

- With the increasing opioid addiction epidemic in our country, effective treatment options are necessary to reverse the trend. Many different pharmacological options are available for opioid maintenance therapy, but research is needed to show which medication has the best efficacy taking into factor such aspects as relapse rates and patient compliance.

## Research Question

- Of the three pharmacological opioid maintenance therapy options endorsed by the Substance Abuse and Mental Health Services administration: methadone, buprenorphine, and naltrexone, which has the best outcomes for patients taking into account efficacy, retention rate, availability, and patient preference?

## Literature Review

A review of the literature provided the following main points:

- Methadone
  - Schukit (2016) states that methadone is superior to placebo for both managing withdrawal and retention in treatment plans.
  - Per Uebelaker et al. (2015), patients do not feel that methadone is the best treatment option, noting barriers such as location and time. They do not feel they are drug free.
- Buprenorphine
  - Nielsen et al. (2016) found that buprenorphine and methadone have similar retention and efficacy, but that buprenorphine may be safer as it has less sedation and respiratory depression.
  - Lobmaier et al. (2010) discussed the abuse potential of buprenorphine and as such recommend the combination of buprenorphine and naloxone.
  - Rosenthal (2013) researched an implantable form of buprenorphine that was found to be three times as effective as placebo and similar to sublingual buprenorphine.
- Naltrexone
  - Lobmaier et al. (2010) found that oral naltrexone is similar only to placebo except in highly motivated individuals for treatment.
  - Nunes et al. (2015) researched an extended release form that has significantly better outcomes than placebo and oral. At 24 weeks, 80% of patients had drug free urines as compared to 15% of patient on oral.
  - Dakwar and Kleber (2015) proposed that the injectable naltrexone can be used to discontinue buprenorphine. For the six patients in the study, only one injection was needed to discontinue use.

## Discussion

Methadone has been used the longest for opioid maintenance treatment. It is tried and true, but patient preference is poor. Patients are limited by location of methadone clinics and the daily dosages.

Buprenorphine is a newer medication, but has shown similar retention and efficacy to methadone. Providers are still limited by certification, but not location so are more abundant. Dosage is not daily.

Naltrexone is the newest treatment of the three. It is little better than placebo as an oral medication, but has shown promise in extended release and implantable forms. These forms allow for at least a month of treatment per dose and have also been used to discontinue the use of buprenorphine. There is also no limitations to prescribers.

Naltrexone allows for more patient autonomy and reduces the time and travel barriers seen with buprenorphine and methadone. It can bring opioid treatment to rural areas that otherwise were without.

Medications for Rehabilitation from an Opioid-Use Disorder, According to the Patient’s Treatment Goal.			
Stage or Function	Full Abstinence from Opioids	Opioid Maintenance	
	Naltrexone	Methadone	Buprenorphine
Action	Blocks opioid high	Long-term maintenance with the use of an oral, long-acting opioid	Long-term maintenance with the use of an oral, long-acting opioid
Restriction	Patient must be opioid-free	No misuse of depressant drugs or medical contraindications; can be used only in specialized programs, not in office-based practices	No misuse of depressant drugs or medical contraindications; can be used in offices of physicians with special training

Schukit, M. A., M.D. (2016, July 2). Treatment of Opioid-Use Disorders. The New England Journal of Medicine, 375(4), 357-368. Retrieved September 26, 2016, from Pubmed

## Applicability to Clinical Practice

First and foremost, there must be a change in how opioids are prescribed by providers. CDC came out with revised guidelines in March of 2016 for primary care providers addressing when to prescribe, how to select, and how to assess and address risk with patients.

Treatment options must also change. Patient preference affects treatment and preference has been towards long acting

medications that require little travel and little time constraints. Drugs like methadone need to be more readily available and have more effective delivery methods.

A third shift must be in attitudes towards opioid misuse. It is a chronic disease similar to hypertension and diabetes. Patients and providers alike must realize that like any other chronic disease, treatment may be on going.

## References

•Dakwar, E., & Kleber, H. D. (2015). Naltrexone-facilitated buprenorphine discontinuation: A feasibility trial. Journal Of Substance Abuse Treatment, 5360-63. doi:10.1016/j.jsat.2015.01.004

•Lembke, A., MD, Humphreys, K., PhD, & Newmark, J., MD. (2016). Weighing the Risks and Benefits of Chronic Opioid Therapy. American Family Physician, 93(12), 982-990. Retrieved September 26, 2016, from Pubmed.

•Lobmaier, P., Gossop, M., Waal, H., & Bramness, J. (2010). The pharmacological treatment of opioid addiction—a clinical perspective. European Journal of Clinical Pharmacology, 66(6), 537-545. doi:10.1007/s00228-010-0793-6

•Nielsen, S., Larance, B., Degenhardt, L., Gowing, L., Kehler, C., & Lintzeris, N. (2016). Opioid agonist treatment for pharmaceutical opioid dependent people. Cochrane Database of Systematic Reviews Reviews. doi:10.1002/14651858.cd011117.pub2

•Nunes, E. V., Krupitsky, E., Ling, W., Zummo, J., Memisoglu, A., Silverman, B. L., & Gastfriend, D. R. (2015). Treating opioid dependence with injectable extended-release naltrexone (XR-NTX): Who will respond?. Journal Of Addiction Medicine, 9(3), 238-243. doi:10.1097/ADM.0000000000000125

•Schukit, M. A., M.D. (2016, July 2). Treatment of Opioid-Use Disorders. The New England Journal of Medicine, 375(4), 357-368. Retrieved September 26, 2016, from Pubmed.

•Uebelacker, L. A., Bailey, G., Herman, D., Anderson, B., & Stein, M. (2016). Patients' Beliefs About Medications are Associated with Stated Preference for Methadone, Buprenorphine, Naltrexone, or no Medication-Assisted Therapy Following Inpatient Opioid Detoxification. Journal of Substance Abuse Treatment, 66, 48-53. doi:10.1016/j.jsat.2016.02.009

## Acknowledgements

I would like to thank UND faculty for their help and encouragement during the process of this project, in particular my advisor Russ Kauffman. Thank yous also go out to both Marilyn Klug for her guidance with the studies chosen and to Erik Christenson for the insight he provided from a pharmacist’s point of view. To my peer group, I thank Savannah Prodzinski, Aggie Ottem, and Angie Shuster for their time, insight, and reassurance.