11-13-2018

Response to criticisms of the 20-m shuttle run test – deflections, distortions, and distractions

Grant R. Tomkinson

University of North Dakota, grant.tomkinson@und.edu

Justin J. Lang

Luc A. Léger

Francisco B. Ortega

Jonatan R. Ruiz

See next page for additional authors

[How does access to this work benefit you? Let us know!](#)

Follow this and additional works at: <https://commons.und.edu/ehb-fac>

Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Grant R. Tomkinson, Justin J. Lang, Luc A. Léger, et al.. "Response to criticisms of the 20-m shuttle run test – deflections, distortions, and distractions" (2018). *Education, Health & Behavior Studies Faculty Publications*. 62.

<https://commons.und.edu/ehb-fac/62>

This Article is brought to you for free and open access by the Department of Education, Health & Behavior Studies at UND Scholarly Commons. It has been accepted for inclusion in Education, Health & Behavior Studies Faculty Publications by an authorized administrator of UND Scholarly Commons. For more information, please contact und.commonson@library.und.edu.

Authors

Grant R. Tomkinson, Justin J. Lang, Luc A. Léger, Francisco B. Ortega, Jonatan R. Ruiz, and Mark S. Tremblay

Title

Response to criticisms of the 20-m shuttle run test — deflections, distortions, and distractions

Authors

Grant R. TOMKINSON,^{*1,2} Justin J. LANG,³ Luc A. LÉGER,⁴ Timothy S. OLDS,^{2,5} Francisco B. ORTEGA,^{6,7} Jonatan R. RUIZ,^{6,7} and Mark S. TREMBLAY.⁸

Institutional affiliations

¹Department of Education, Health and Behavior Studies, University of North Dakota, Grand Forks, USA

²Alliance for Research in Exercise, Nutrition and Activity (ARENA), School of Health Sciences & Sansom Institute for Health Research, University of South Australia, Adelaide, Australia

³Centre for Surveillance and Applied Research, Public Health Agency of Canada, Ottawa, Canada

⁴Department of Kinesiology, University of Montreal, Montreal, Canada

⁵Murdoch Children's Research Institute, Melbourne, Australia

⁶PROmoting FITness and Health through physical activity research group (PROFITH), Department of Physical Education and Sports, Faculty of Sport Sciences, University of Granada, Granada, Spain

⁷Department of Biosciences and Nutrition, Karolinska Institutet, Huddinge, Sweden

⁸Healthy Active Living and Obesity (HALO) Research Group, CHEO Research Institute, Ottawa, Canada

Corresponding author

Professor Grant R. TOMKINSON*

✉ Department of Education, Health and Behavior Studies
University of North Dakota
2751 2nd Avenue North, Stop 8235
Grand Forks, ND, 58202, USA

☎ +1 701-777-4041

📧 grant.tomkinson@und.edu

Keywords: cardiorespiratory fitness; children; 20-m shuttle run; population health.

Word count: 826

In their editorial, Armstrong and Welsman (1) suggest that the 20-m shuttle run test (20mSRT) (mis)represents and (mis)interprets youth cardiorespiratory fitness (CRF), and potentially (mis)informs health promotion and clinical practice. Their main arguments are: (a) the 20mSRT only provides an estimate of CRF (i.e., peak $\dot{V}O_2$), and (b) estimates are ratio-scaled (i.e., expressed relative to body mass). In this response we provide several reasons, rooted in evidence, which refute their interpretation of our work.

CRF measures the body's capacity to deliver and utilise oxygen for energy transfer to support muscle activity during physical activity (2). The 20mSRT provides a simple, single measure that assesses the integrated responses of the physiological systems' ability to perform progressive aerobic exercise. Unfortunately, it does not provide specific information on the function or contribution of specific systems that can be obtained from a gas analysed peak $\dot{V}O_2$ test. The 20mSRT is a good measure of functional exercise capacity that authentically imitates youth physical activities (e.g., running, agility) in a natural setting. At the individual level, the 20mSRT is a true indicator of peak $\dot{V}O_2$ (absolute or relative to body mass) because peak $\dot{V}O_2$ is achieved at the end of a maximal performance (3). While it is widely accepted that gas-analysed peak $\dot{V}O_2$ is the criterion physiological measure of CRF, other factors also contribute to CRF (e.g., mechanical efficiency, $\dot{V}O_2$ kinetics etc.). 20mSRT performance, as well as treadmill running/walking, is also affected by physical (e.g., excess adiposity) and psychosocial (e.g., self-efficacy, motivation) factors.

Evidence supports the use of the 20mSRT in childhood and adolescence as a powerful marker of current health status (4), independent of adiposity (5), and a predictor of future health (4), making it very useful for population health research. Analogously, BMI is a widely used population health measure with moderate criterion validity and is more strongly associated with health and mortality than criterion-measured adiposity (e.g., hydrostatic weighing) (6). Like BMI, the 20mSRT could be a useful population health measure to identify at-risk populations (4).

Unlike gas-analysed peak $\dot{V}O_2$, the 20mSRT has strong utility and scalability for population health surveillance (4). A recent systematic review of field-based CRF tests identified the 20mSRT as the most scalable for school-based testing (7). It is routinely used as a preferred option to monitor progress in response to physical activity programs because of the low cost of equipment, and its ability to test large groups of participants, simultaneously. For these reasons, the 20mSRT is scalable for international research, especially for data collection in low-income countries.

Armstrong and Welsman (1) highlighted that an earlier study (2) reported a very small trend in the gas-analysed peak $\dot{V}O_2$ (mL/kg/min) of ~4000 youth from five countries between 1962 and 1994. Differences in testing modality/protocols/conditions, metabolic systems, and maximum effort criteria affect the interpretation of such trends. Although now out-dated, the reported trend probably reflects no meaningful change in peak $\dot{V}O_2$ in a non-representative sample of volunteers who likely had an athletic predisposition (2). In contrast, our data of ~1 million youth from 19 countries, which likely provide a better representation of population trends in CRF,

indicate a decline in 20mSRT performance between 1981 and 2014, with no recent change in several countries (4). Available data also indicate that declines in 20mSRT performance are independent of changes in adiposity, suggesting corresponding declines in habitual physical activity levels (8).

A meta-analysis reported the 20mSRT's criterion validity as moderate for estimating peak $\dot{V}O_2$ in youth (9), which compares favourably to other field-based CRF tests (e.g., distance/timed runs). We agree with Armstrong and Welsman (1) that allometric scaling can help correct the limitations of traditional ratio-scaled peak $\dot{V}O_2$ which fails to produce peak $\dot{V}O_2$ independent of body mass. Unfortunately, even they have reported body mass exponents ranging from 0.37 to 0.94, highlighting that allometric exponents are sample-specific (10). The continued use of ratio-scaled peak $\dot{V}O_2$ is (in their words) “in the absence of a universally applicable alternative” (10). Despite this, we have recommended reporting 20mSRT results in the measured units (e.g., number of laps, stages, peak running speed), in addition to estimated peak $\dot{V}O_2$, when necessary, with a caveat that the latter results in prediction error. Often at the request of reviewers/editors, we have estimated peak $\dot{V}O_2$ when pooling 20mSRT data to report international norms, trends, and health-related criterion-referenced standards. Regardless, when comparing the CRF of youth in weight-bearing activities and for better transfer to activities of daily living, expressing peak $\dot{V}O_2$ per unit body mass is appropriate.

Armstrong and Welsman (1) suggest that we have (mis)represented and (mis)interpreted youth CRF while potentially (mis)informing health promotion and clinical practice. We strongly disagree. Lab-based peak $\dot{V}O_2$ is an excellent prognostic measure at the individual level. In

contrast, the 20mSRT has good feasibility, utility, and scalability for population health surveillance to monitor trends, and inform policymaking and public health planning. Despite their deflections, distortions, and distractions, the 20mSRT is pragmatic, easily understood and routinely used internationally to measure CRF, and we hope this continues.

Acknowledgements

None

Contributors

GRT and JLL drafted the editorial. All authors reviewed and approved the final version.

Competing interests

None declared

Funding information

None

Disclaimer

The content and views expressed in this article are those of the authors and do not necessarily reflect those of the Government of Canada.

References

- 1 Armstrong N, Welsman J. The 20-m shuttle run — (mis)representation, (mis)interpretation, and (mis)use. *Br J Sports Med* 2018; in press.
- 2 Armstrong N, Tomkinson G, Ekelund U. Aerobic fitness and its relationship to sport, exercise training and habitual physical activity during youth. *Br J Sports Med* 2011;45(11):849–858.
- 3 Ruiz JR, Silva G, Oliveira N, et al. Criterion-related validity of the 20-m shuttle run test in youths aged 13–19 years. *J Sports Sci* 2009;27(9):899–906.
- 4 Lang JJ, Tomkinson GR, Janssen I, et al. Making a case for cardiorespiratory fitness surveillance among children and youth. *Exerc Sport Sci Rev* 2018;46(2):66–75.
- 5 Machado-Rodrigues AM, Leite N, Coelho-E-Silva MJ, et al. Independent association of clustered risk factors with cardiorespiratory fitness in youth aged 11–17 years. *Ann Hum Biol* 2014;41(3):271–276.
- 6 Ortega FB, Sui X, Lavie CJ, et al. Body mass index, the most widely used but also widely criticized index: would a criterion standard measure of total body fat be a better predictor of cardiovascular disease mortality? *Mayo Clin Proc.* 2016;91(4):443–455.
- 7 Domone S, Mann S, Sandercock G, et al. A method by which to assess the scalability of field-based fitness tests of cardiorespiratory fitness among schoolchildren. *Sports Med* 2016;46(12):1819–1831.
- 8 Sandercock GR, Ogunleye A, Voss C. Six-year changes in body mass index and cardiorespiratory fitness of English schoolchildren from an affluent area. *Int J Obes (Lond)* 2015;39(10):1504–1507.

- 9 Mayorga-Vega D, Aguilar-Soto P, Viciano J. Criterion-related validity of the 20-m shuttle run test for estimating cardiorespiratory fitness: A meta-analysis. *J Sports Sci Med* 2015;14(3):536–547.
- 10 Welsman J, Armstrong N. Interpreting aerobic fitness in youth: the fallacy of ratio scaling. *Pediatr Exerc Sci* 2018:1–7. doi: 10.1123/pes.2018-0141.